

MCM3 antibody - C-terminal region

Rabbit Polyclonal Antibody

Catalog # AI11009

Product Information

Application	WB, IHC
Primary Accession	P25205
Other Accession	NM_002388 , NP_002379
Reactivity	Human, Mouse, Rat, Rabbit, Dog, Horse, Bovine, Yeast
Predicted	Human, Mouse, Rat, Chicken, Horse, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	90981

Additional Information

Gene ID	4172
Alias Symbol	HCC5, MGC1157, P1-MCM3, P1.h, RLFB
Other Names	DNA replication licensing factor MCM3, 3.6.4.12, DNA polymerase alpha holoenzyme-associated protein P1, P1-MCM3, RLF subunit beta, p102, MCM3
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 50 ul of distilled water. Final anti-MCM3 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
Precautions	MCM3 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	MCM3 (HGNC:6945)
Function	Acts as a component of the MCM2-7 complex (MCM complex) which is the replicative helicase essential for 'once per cell cycle' DNA replication initiation and elongation in eukaryotic cells. Core component of CDC45-MCM-GINS (CMG) helicase, the molecular machine that unwinds template DNA during replication, and around which the replisome is built (PubMed: 32453425 , PubMed: 34694004 , PubMed: 34700328 , PubMed: 35585232). The active ATPase sites in the MCM2-7 ring are formed through the interaction surfaces of two neighboring subunits such that a critical structure of a conserved arginine finger motif is provided in trans relative to the ATP-binding site of the Walker A box of the adjacent subunit. The six ATPase active sites, however, are likely

to contribute differentially to the complex helicase activity (PubMed:[32453425](#)). Required for the entry in S phase and for cell division (Probable).


Cellular Location

Nucleus. Chromosome. Note=Associated with chromatin before the formation of nuclei and detaches from it as DNA replication progresses.

References

Lin,D.I., (2008) Proc. Natl. Acad. Sci. U.S.A. 105 (23), 8079-8084 Reconstitution and Storage:For short term use, store at 2-8C up to 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.

Images



WB Suggested Anti-MCM3 Antibody Titration: 0.2-1 µg/ml
ELISA Titer: 1:312500
Positive Control: 721_B cell lysate
MCM3 is supported by BioGPS gene expression data to be expressed in 721_B

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.