

Acid antibody - C-terminal region

Rabbit Polyclonal Antibody

Catalog # AI11035

Product Information

Application	WB
Primary Accession	Q5EE38
Other Accession	NM_001012638 , NP_001012656
Reactivity	Human, Mouse, Rat, Dog, Horse, Bovine
Predicted	Human, Mouse, Rat, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	44713

Additional Information


Gene ID	497652
Other Names	Adrenocortical dysplasia protein, Acid
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 50 ul of distilled water. Final anti-Acid antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
Precautions	Acid antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	Acid {ECO:0000312 MGI:MGI:87873}
Function	Component of the shelterin complex (telosome) that is involved in the regulation of telomere length and protection. Shelterin associates with arrays of double-stranded TTAGGG repeats added by telomerase and protects chromosome ends. Without its protective activity, telomeres are no longer hidden from the DNA damage surveillance and chromosome ends are inappropriately processed by DNA repair pathways. Promotes binding of POT1 to single-stranded telomeric DNA. Modulates the inhibitory effects of POT1 on telomere elongation. The ACD-POT1 heterodimer enhances telomere elongation by recruiting telomerase to telomeres and increasing its processivity (By similarity). May play a role in organogenesis (PubMed: 15537664).

Cellular Location	Nucleus {ECO:0000250 UniProtKB:Q96AP0}. Chromosome, telomere {ECO:0000250 UniProtKB:Q96AP0}
Tissue Location	Ubiquitous..

Images



WB Suggested Anti-Acd Antibody Titration: 0.2-1 µg/ml
Positive Control: Mouse Muscle

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.