

Foxa3 antibody - C-terminal region

Rabbit Polyclonal Antibody Catalog # AI11057

Product Information

<u>84</u>
<u>008260</u> , <u>NP_032286</u>
an, Mouse, Rat, Pig, Dog, Horse, Bovine
an, Mouse, Rat, Dog, Horse, Bovine
it
lonal
1

Additional Information

Gene ID	15377
Alias Symbol Other Names	Hnf-3g, Hnf3g, Tcf-3g, Tcf3g Hepatocyte nuclear factor 3-gamma, HNF-3-gamma, HNF-3G, Forkhead box protein A3, Foxa3, Hnf3g, Tcf-3g, Tcf3g
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 50 ul of distilled water. Final anti-Foxa3 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
Precautions	Foxa3 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	Foxa3
Synonyms	Hnf3g, Tcf-3g, Tcf3g
Function	Transcription factor that is thought to act as a 'pioneer' factor opening the compacted chromatin for other proteins through interactions with nucleosomal core histones and thereby replacing linker histones at target enhancer and/or promoter sites (By similarity). Originally described as a transcription activator for a number of liver genes such as AFP, albumin, tyrosine aminotransferase, PEPCK, etc. Interacts with the cis-acting regulatory regions of these genes. Involved in glucose homeostasis; activates GLUT2 transcription.

	Involved in regulation of spermatogenesis; required for the maintenance of the testicular germ cell population and male fertility.
Cellular Location	Nucleus.
Tissue Location	Restricted mainly to endoderm-derived tissues (lung, liver, stomach, and small intestine), also present additionally in ovary, testis, heart, and adipose tissue, but missing from lung
Images	
90 kDa_ 65 kDa_ 40 kDa_ 31 kDa_ 22 kDa_	WB Suggested Anti-Foxa3 Antibody Titration: 0.2-1 μg/ml ELISA Titer: 1:312500 Positive Control: Mouse Brain

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.