

NFATC2 antibody - N-terminal region

Rabbit Polyclonal Antibody

Catalog # AI11130

Product Information

Application	WB
Primary Accession	Q60591-3
Other Accession	NM_010899 , NP_035029
Reactivity	Human, Mouse, Rat, Pig, Dog, Horse, Bovine
Predicted	Human, Mouse, Rat, Pig, Dog, Horse, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	102 KDa

Additional Information

Alias Symbol	NFAT1, Nfatp, NF-ATc2, NFAT1-D, AI607462
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 100 ul of distilled water. Final anti-NFATC2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
Precautions	NFATC2 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.


Protein Information

References

Glud,S.Z., et al., (2005) Blood 106 (10), 3546-3552
Reconstitution and Storage:For short term use, store at 2-8C up to 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.
Publications:Heer, R. et al. Phenotypic modulation of human urinary tract stroma-derived fibroblasts by transforming growth factor beta3. Urology 76, 509.e13-20 (2010). WB, IP, Mouse, Human, Rat, H, Bovine, Dog, Pig20546875
Nagamoto-Combs, K. & Combs, C. K. Microglial phenotype is regulated by activity of the transcription factor, NFAT (nuclear factor of activated T cells). J. Neurosci. 30, 9641-6 (2010). WB, IP, Mouse, Human, Rat, H, Bovine, Dog, Pig20631193

Images

WB Suggested Anti-NFATC2 Antibody Titration: 1.25µg/ml



ELISA Titer: 1:312500
Positive Control: SP2/0 cell lysate

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.