

Rorc antibody - N-terminal region

Rabbit Polyclonal Antibody

Catalog # AI11132

Product Information

Application	WB, IHC
Primary Accession	P51450
Other Accession	NM_011281 , NP_035411
Reactivity	Human, Mouse, Rat, Pig, Dog
Predicted	Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	58117

Additional Information

Gene ID	19885
Alias Symbol Other Names	Nr1f3, RORgamma, TOR, Thor Nuclear receptor ROR-gamma, Nuclear receptor RZR-gamma, Nuclear receptor subfamily 1 group F member 3, RAR-related orphan receptor C, Retinoid-related orphan receptor-gamma, Thymus orphan receptor, TOR, Rorc, Nr1f3, Rorg, Thor
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 50 ul of distilled water. Final anti-Rorc antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
Precautions	Rorc antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	Rorc
Synonyms	Nr1f3, Rorg, Thor
Function	Nuclear receptor that binds DNA as a monomer to ROR response elements (RORE) containing a single core motif half-site 5'-AGGTCA-3' preceded by a short A-T-rich sequence. Key regulator of cellular differentiation, immunity, peripheral circadian rhythm as well as lipid, steroid, xenobiotics and glucose metabolism. Considered to have intrinsic transcriptional activity, have some natural ligands like oxysterols that act as agonists (25-hydroxycholesterol) or

inverse agonists (7-oxygenated sterols), enhancing or repressing the transcriptional activity, respectively. Recruits distinct combinations of cofactors to target gene regulatory regions to modulate their transcriptional expression, depending on the tissue, time and promoter contexts (PubMed:[17666523](#), PubMed:[19381306](#), PubMed:[19965867](#), PubMed:[21853531](#), PubMed:[22789990](#), PubMed:[23723244](#)). Regulates the circadian expression of clock genes such as CRY1, BMAL1 and NR1D1 in peripheral tissues and in a tissue-selective manner (PubMed:[22753030](#)). Competes with NR1D1 for binding to their shared DNA response element on some clock genes such as BMAL1, CRY1 and NR1D1 itself, resulting in NR1D1-mediated repression or RORC-mediated activation of the expression, leading to the circadian pattern of clock genes expression. Therefore influences the period length and stability of the clock (PubMed:[22753030](#)). Involved in the regulation of the rhythmic expression of genes involved in glucose and lipid metabolism, including PLIN2 and AVPR1A. Negative regulator of adipocyte differentiation through the regulation of early phase genes expression, such as MMP3. Controls adipogenesis as well as adipocyte size and modulates insulin sensitivity in obesity. In liver, has specific and redundant functions with RORA as positive or negative modulator of expression of genes encoding phase I and Phase II proteins involved in the metabolism of lipids, steroids and xenobiotics, such as SULT1E1 (PubMed:[21853531](#)). Also plays a role in the regulation of hepatocyte glucose metabolism through the regulation of G6PC1 and PCK1. Regulates the rhythmic expression of PROX1 and promotes its nuclear localization. Plays an indispensable role in the induction of IFN-gamma dependent anti-mycobacterial systemic immunity (By similarity).


Cellular Location

Nucleus {ECO:0000255 | PROSITE-ProRule:PRU00407, ECO:0000269 | PubMed:18368049, ECO:0000269 | PubMed:21853531, ECO:0000269 | PubMed:22753030}

Tissue Location

Expressed in immature Vgamma2 gamma-delta T-cells (at protein level) (PubMed:23562159). Expressed in the liver (PubMed:17666523, PubMed:22753030). Expressed in the heart (PubMed:17666523). Expressed in the kidney, jejunum, and brown adipose tissue (PubMed:22753030). [Isoform 2]: Expressed in the thymus, primarily in immature thymocytes, including Vgamma2 gamma-delta T-cells (at protein levels) (PubMed:10602018, PubMed:9881970). Also expressed in a subset of mature T(H)17 cells (PubMed:18164222). Not expressed in the spleen or bone marrow (PubMed:9881970).

Images



WB Suggested Anti-Rorc Antibody Titration: 1 µg/ml
Positive Control: Mouse Heart lysate

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.