

Jarid2 antibody - middle region

Rabbit Polyclonal Antibody

Catalog # AI11180

Product Information

Application	WB
Primary Accession	Q62315
Other Accession	NM_021878 , NP_068678
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Dog
Predicted	Mouse, Rabbit, Pig
Host	Rabbit
Clonality	Polyclonal
Calculated MW	137445

Additional Information

Gene ID	16468
Alias Symbol	AI317256, AU045941, C79929, C79931, Jmj, jumonji
Other Names	Protein Jumonji, Jumonji/ARID domain-containing protein 2, Jarid2, Jmj
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 50 ul of distilled water. Final anti-Jarid2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
Precautions	Jarid2 antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	Jarid2
Synonyms	Jmj
Function	Regulator of histone methyltransferase complexes that plays an essential role in embryonic development, including heart and liver development, neural tube fusion process and hematopoiesis (PubMed: 10807864 , PubMed: 12852854 , PubMed: 12890668 , PubMed: 15542826 , PubMed: 15870077 , PubMed: 19010785 , PubMed: 20064375 , PubMed: 20064376 , PubMed: 20075857). Acts as an accessory subunit for the core PRC2 (Polycomb repressive complex 2) complex, which mediates histone H3K27 (H3K27me3) trimethylation on chromatin (PubMed: 20064375 , PubMed: 20064376). Binds DNA and mediates the recruitment of the PRC2

complex to target genes in embryonic stem cells, thereby playing a key role in stem cell differentiation and normal embryonic development (PubMed:[20064375](#), PubMed:[20075857](#)). In cardiac cells, it is required to repress expression of cyclin-D1 (CCND1) by activating methylation of 'Lys-9' of histone H3 (H3K9me) by the GLP1/EHMT1 and G9a/EHMT2 histone methyltransferases (PubMed:[12852854](#), PubMed:[12890668](#), PubMed:[19010785](#)). Also acts as a transcriptional repressor of ANF via its interaction with GATA4 and NKX2-5 (PubMed:[15542826](#)). Participates in the negative regulation of cell proliferation signaling (PubMed:[10913339](#)). Does not have histone demethylase activity (PubMed:[20064376](#)).


Cellular Location

Nucleus {ECO:0000255 | PROSITE-ProRule:PRU00355, ECO:0000255 | PROSITE-ProRule:PRU00537, ECO:0000269 | PubMed:10807864, ECO:0000269 | PubMed:10913339, ECO:0000269 | PubMed:20064375, ECO:0000269 | PubMed:20064376}. Note=Colocalizes with the PRC2 complex on chromatin

Tissue Location

Widely expressed in embryos. In adults, expressed at high levels in heart, skeletal muscle, brain and thymus

Images



WB Suggested Anti-Jarid2 Antibody Titration: 0.2-1 µg/ml
ELISA Titer: 1:1562500
Positive Control: Mouse Liver

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.