

Vgll2 antibody - C-terminal region

Rabbit Polyclonal Antibody Catalog # AI11240

Product Information

Application WB

Primary Accession Q8BGW8

Other Accession <u>NM 153786, NP 722481</u>

Reactivity Human, Mouse, Rat, Rabbit, Dog

Predicted Human, Mouse, Rat, Rabbit, Chicken, Dog

Host Rabbit
Clonality Polyclonal
Calculated MW 34064

Additional Information

Gene ID 215031

Alias Symbol C130057C21Rik, VITO-1, Vito1

Other Names Transcription cofactor vestigial-like protein 2, Vgl-2, Protein VITO1, Vgll2, Vgl2,

Vito1

Format Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium

azide and 2% sucrose.

Reconstitution & Storage Add 50 ul of distilled water. Final anti-Vgll2 antibody concentration is 1 mg/ml

in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C.

Avoid repeat freeze-thaw cycles.

Precautions Vgll2 antibody - C-terminal region is for research use only and not for use in

diagnostic or therapeutic procedures.

Protein Information

Name VgII2

Synonyms Vgl2, Vito1

Function May act as a specific coactivator for the mammalian TEFs. May play a role in

the development of skeletal muscles.

Cellular Location Nucleus.

Tissue Location Skeletal muscle specific.

Images

90 kDa_ 65 kDa_ 40 kDa_ 31 kDa_ 22 kDa_

WB Suggested Anti-Vgll2 Antibody Titration: 0.2-1 $\mu g/ml$

ELISA Titer: 1:312500

Positive Control: SP2/0 cell lysate

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.