

# G3BP antibody - N-terminal region

Rabbit Polyclonal Antibody

Catalog # AI11302

## Product Information

Application	WB
Primary Accession	<a href="#">Q13283</a>
Other Accession	<a href="#">NM_005754</a> , <a href="#">NP_005745</a>
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Dog, Horse, Bovine
Predicted	Human, Dog, Horse, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	52164

## Additional Information

Gene ID	10146
Alias Symbol	G3BP, HDH-VIII
Other Names	Ras GTPase-activating protein-binding protein 1, G3BP-1, 3.6.4.12, 3.6.4.13, ATP-dependent DNA helicase VIII, hDH VIII, GAP SH3 domain-binding protein 1, G3BP1, G3BP
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 100 ul of distilled water. Final anti-G3BP antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
Precautions	G3BP antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

Name	G3BP1 {ECO:0000303 PubMed:23279204, ECO:0000312 HGNC:HGNC:30292}
Function	Protein involved in various processes, such as stress granule formation and innate immunity (PubMed: <a href="#">12642610</a> , PubMed: <a href="#">20180778</a> , PubMed: <a href="#">23279204</a> , PubMed: <a href="#">30510222</a> , PubMed: <a href="#">30804210</a> ). Plays an essential role in stress granule formation (PubMed: <a href="#">12642610</a> , PubMed: <a href="#">20180778</a> , PubMed: <a href="#">23279204</a> , PubMed: <a href="#">32302570</a> , PubMed: <a href="#">32302571</a> , PubMed: <a href="#">32302572</a> , PubMed: <a href="#">34739333</a> , PubMed: <a href="#">35977029</a> , PubMed: <a href="#">36183834</a> , PubMed: <a href="#">36279435</a> , PubMed: <a href="#">36692217</a> , PubMed: <a href="#">37379838</a> ). Stress granules are membraneless compartments that

store mRNAs and proteins, such as stalled translation pre-initiation complexes, in response to stress (PubMed:[12642610](#), PubMed:[20180778](#), PubMed:[23279204](#), PubMed:[27022092](#), PubMed:[32302570](#), PubMed:[32302571](#), PubMed:[32302572](#), PubMed:[36279435](#), PubMed:[37379838](#)). Promotes formation of stress granules phase-separated membraneless compartment by undergoing liquid-liquid phase separation (LLPS) upon unfolded RNA-binding: functions as a molecular switch that triggers RNA-dependent LLPS in response to a rise in intracellular free RNA concentrations (PubMed:[32302570](#), PubMed:[32302571](#), PubMed:[32302572](#), PubMed:[34739333](#), PubMed:[36279435](#), PubMed:[36692217](#)). Also acts as an ATP- and magnesium-dependent helicase: unwinds DNA/DNA, RNA/DNA, and RNA/RNA substrates with comparable efficiency (PubMed:[9889278](#)). Acts unidirectionally by moving in the 5' to 3' direction along the bound single-stranded DNA (PubMed:[9889278](#)). Unwinds preferentially partial DNA and RNA duplexes having a 17 bp annealed portion and either a hanging 3' tail or hanging tails at both 5'- and 3'-ends (PubMed:[9889278](#)). Plays an essential role in innate immunity by promoting CGAS and RIGI activity (PubMed:[30510222](#), PubMed:[30804210](#)). Participates in the DNA-triggered cGAS/STING pathway by promoting the DNA binding and activation of CGAS (PubMed:[30510222](#)). Triggers the condensation of cGAS, a process probably linked to the formation of membrane-less organelles (PubMed:[34779554](#)). Also enhances RIGI-induced type I interferon production probably by helping RIGI at sensing pathogenic RNA (PubMed:[30804210](#)). May also act as a phosphorylation- dependent sequence-specific endoribonuclease in vitro: Cleaves exclusively between cytosine and adenine and cleaves MYC mRNA preferentially at the 3'-UTR (PubMed:[11604510](#)).

## Cellular Location

Cytoplasm, cytosol. Perikaryon {ECO:0000250|UniProtKB:P97855}. Cytoplasm, Stress granule. Nucleus Note=Cytoplasmic in proliferating cells (PubMed:[11604510](#)). Cytosolic and partially nuclear in resting cells (PubMed:[11604510](#)). Recruited to stress granules in response to arsenite treatment (PubMed:[12642610](#), PubMed:[20180778](#)). The unphosphorylated form is recruited to stress granules (PubMed:[12642610](#)). HRAS signaling contributes to this process by regulating G3BP dephosphorylation (PubMed:[12642610](#))


## Tissue Location

Ubiquitous..

## References

Kociok,N., et al., (1999) J. Cell. Biochem. 74 (2), 194-201  
Reconstitution and Storage:For short term use, store at 2-8C up to 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.

## Images



WB Suggested Anti-G3BP Antibody Titration: 2.5µg/ml  
ELISA Titer: 1:312500  
Positive Control: HepG2 cell lysate