

G3BP antibody - N-terminal region

Rabbit Polyclonal Antibody
Catalog # AI11302

Product Information

Application	WB
Primary Accession	Q13283
Other Accession	NM_005754 , NP_005745
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Dog, Horse, Bovine
Predicted	Human, Dog, Horse, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	52164

Additional Information

Gene ID	10146
Alias Symbol	G3BP, HDH-VIII
Other Names	Ras GTPase-activating protein-binding protein 1, G3BP-1, 3.6.4.12, 3.6.4.13, ATP-dependent DNA helicase VIII, hDH VIII, GAP SH3 domain-binding protein 1, G3BP1, G3BP
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 100 ul of distilled water. Final anti-G3BP antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
Precautions	G3BP antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	G3BP1 {ECO:0000303 PubMed:23279204, ECO:0000312 HGNC:HGNC:30292}
Function	Protein involved in various processes, such as stress granule formation and innate immunity (PubMed: 12642610 , PubMed: 20180778 , PubMed: 23279204 , PubMed: 30510222 , PubMed: 30804210). Plays an essential role in stress granule formation (PubMed: 12642610 , PubMed: 20180778 , PubMed: 23279204 , PubMed: 32302570 , PubMed: 32302571 , PubMed: 32302572 , PubMed: 34739333 , PubMed: 35977029 , PubMed: 36183834 , PubMed: 36279435 , PubMed: 36692217 , PubMed: 37379838). Stress granules are membraneless compartments that

store mRNAs and proteins, such as stalled translation pre-initiation complexes, in response to stress (PubMed:[12642610](#), PubMed:[20180778](#), PubMed:[23279204](#), PubMed:[27022092](#), PubMed:[32302570](#), PubMed:[32302571](#), PubMed:[32302572](#), PubMed:[36279435](#), PubMed:[37379838](#)). Promotes formation of stress granules phase-separated membraneless compartment by undergoing liquid-liquid phase separation (LLPS) upon unfolded RNA-binding: functions as a molecular switch that triggers RNA-dependent LLPS in response to a rise in intracellular free RNA concentrations (PubMed:[32302570](#), PubMed:[32302571](#), PubMed:[32302572](#), PubMed:[34739333](#), PubMed:[36279435](#), PubMed:[36692217](#)). Also acts as an ATP- and magnesium-dependent helicase: unwinds DNA/DNA, RNA/DNA, and RNA/RNA substrates with comparable efficiency (PubMed:[9889278](#)). Acts unidirectionally by moving in the 5' to 3' direction along the bound single-stranded DNA (PubMed:[9889278](#)). Unwinds preferentially partial DNA and RNA duplexes having a 17 bp annealed portion and either a hanging 3' tail or hanging tails at both 5'- and 3'-ends (PubMed:[9889278](#)). Plays an essential role in innate immunity by promoting CGAS and RIGI activity (PubMed:[30510222](#), PubMed:[30804210](#)). Participates in the DNA-triggered cGAS/STING pathway by promoting the DNA binding and activation of CGAS (PubMed:[30510222](#)). Triggers the condensation of cGAS, a process probably linked to the formation of membrane-less organelles (PubMed:[34779554](#)). Also enhances RIGI-induced type I interferon production probably by helping RIGI at sensing pathogenic RNA (PubMed:[30804210](#)). May also act as a phosphorylation- dependent sequence-specific endoribonuclease in vitro: Cleaves exclusively between cytosine and adenine and cleaves MYC mRNA preferentially at the 3'-UTR (PubMed:[11604510](#)).

Cellular Location

Cytoplasm, cytosol. Perikaryon {ECO:0000250|UniProtKB:P97855}. Cytoplasm, Stress granule. Nucleus Note=Cytoplasmic in proliferating cells (PubMed:[11604510](#)). Cytosolic and partially nuclear in resting cells (PubMed:[11604510](#)). Recruited to stress granules in response to arsenite treatment (PubMed:[12642610](#), PubMed:[20180778](#)). The unphosphorylated form is recruited to stress granules (PubMed:[12642610](#)). HRAS signaling contributes to this process by regulating G3BP dephosphorylation (PubMed:[12642610](#))


Tissue Location

Ubiquitous..

References

Kociok,N., et al., (1999) J. Cell. Biochem. 74 (2), 194-201
Reconstitution and Storage:For short term use, store at 2-8C up to 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.

Images



WB Suggested Anti-G3BP Antibody Titration: 2.5µg/ml
ELISA Titer: 1:312500
Positive Control: HepG2 cell lysate