

# BLOC1S1 Antibody - N-terminal region

Rabbit Polyclonal Antibody

Catalog # AI11354

## Product Information

<b>Application</b>	WB
<b>Primary Accession</b>	<a href="#">P78537</a>
<b>Other Accession</b>	<a href="#">NM_001487</a> , <a href="#">NP_001478</a>
<b>Reactivity</b>	Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Dog, Horse, Bovine
<b>Predicted</b>	Human, Mouse, Rat, Rabbit, Zebrafish, Pig
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Calculated MW</b>	17263

## Additional Information

<b>Gene ID</b>	2647
<b>Alias Symbol</b>	BLOS1, GCN5L1, MICoA, RT14
<b>Other Names</b>	Biogenesis of lysosome-related organelles complex 1 subunit 1, BLOC-1 subunit 1, GCN5-like protein 1, Protein RT14, BLOC1S1, BLOS1, GCN5L1, RT14
<b>Format</b>	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
<b>Reconstitution &amp; Storage</b>	Add 50 ul of distilled water. Final anti-BLOC1S1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
<b>Precautions</b>	BLOC1S1 Antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

<b>Name</b>	BLOC1S1
<b>Synonyms</b>	BLOS1, GCN5L1 {ECO:0000303 PubMed:382816}
<b>Function</b>	Component of the BLOC-1 complex, a complex that is required for normal biogenesis of lysosome-related organelles (LRO), such as platelet dense granules and melanosomes (PubMed: <a href="#">17182842</a> ). In concert with the AP-3 complex, the BLOC-1 complex is required to target membrane protein cargos into vesicles assembled at cell bodies for delivery into neurites and nerve terminals (PubMed: <a href="#">17182842</a> ). The BLOC-1 complex, in association with SNARE proteins, is also proposed to be involved in neurite extension (PubMed: <a href="#">17182842</a> ). As part of the BORC complex may play a role in


lysosomes movement and localization at the cell periphery (PubMed:[25898167](#)). Associated with the cytosolic face of lysosomes, the BORC complex may recruit ARL8B and couple lysosomes to microtubule plus-end-directed kinesin motor (PubMed:[25898167](#)).

## Cellular Location

Mitochondrion intermembrane space. Mitochondrion matrix. Cytoplasm, cytosol. Lysosome membrane

## Images

---



Host: Rabbit  
Target Name: BLOC1S1  
Sample Tissue: Lung Tumor lysates  
Antibody Dilution: 1.0µg/ml

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.