

ZFP36 antibody - N-terminal region

Rabbit Polyclonal Antibody

Catalog # AI11388

Product Information

| | |
|-------------------|---|
| Application | WB |
| Primary Accession | P26651 |
| Other Accession | NM_003407 , NP_003398 |
| Reactivity | Human |
| Predicted | Human |
| Host | Rabbit |
| Clonality | Polyclonal |
| Calculated MW | 34003 |

Additional Information

| | |
|-----------------------------|--|
| Gene ID | 7538 |
| Alias Symbol Other Names | TTP, GOS24, GOS24, TIS11, NUP475, zfp-36, RNF162A Tristetraprolin, TTP, G0/G1 switch regulatory protein 24, Growth factor-inducible nuclear protein NUP475, Protein TIS11A, TIS11, Zinc finger protein 36 homolog, Zfp-36, ZFP36, GOS24, RNF162A, TIS11A, TTP |
| Format | Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose. |
| Reconstitution & Storage | Add 50 ul of distilled water. Final anti-ZFP36 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles. |
| Precautions | ZFP36 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures. |

Protein Information

| | |
|----------|---|
| Name | ZFP36 (HGNC:12862) |
| Function | Zinc-finger RNA-binding protein that destabilizes several cytoplasmic AU-rich element (ARE)-containing mRNA transcripts by promoting their poly(A) tail removal or deadenylation, and hence provide a mechanism for attenuating protein synthesis (PubMed: 10330172 , PubMed: 10751406 , PubMed: 11279239 , PubMed: 12115244 , PubMed: 12748283 , PubMed: 15187101 , PubMed: 15634918 , PubMed: 16702957 , PubMed: 17030620 , PubMed: 20221403 , PubMed: 20702587 , PubMed: 21775632 , PubMed: 23644599 , PubMed: 25815583 , PubMed: 27193233 , PubMed: 31439631 , PubMed: 9703499). Acts as an 3'-untranslated region (UTR) |

ARE mRNA-binding adapter protein to communicate signaling events to the mRNA decay machinery (PubMed:[15687258](#), PubMed:[23644599](#)). Recruits deadenylase CNOT7 (and probably the CCR4-NOT complex) via association with CNOT1, and hence promotes ARE-mediated mRNA deadenylation (PubMed:[23644599](#)). Functions also by recruiting components of the cytoplasmic RNA decay machinery to the bound ARE-containing mRNAs (PubMed:[11719186](#), PubMed:[12748283](#), PubMed:[15687258](#), PubMed:[16364915](#)). Self regulates by destabilizing its own mRNA (PubMed:[15187101](#)). Binds to 3'-UTR ARE of numerous mRNAs and of its own mRNA (PubMed:[10330172](#), PubMed:[10751406](#), PubMed:[12115244](#), PubMed:[15187101](#), PubMed:[15634918](#), PubMed:[16702957](#), PubMed:[17030620](#), PubMed:[19188452](#), PubMed:[20221403](#), PubMed:[20702587](#), PubMed:[21775632](#), PubMed:[25815583](#)). Plays a role in anti-inflammatory responses; suppresses tumor necrosis factor (TNF)-alpha production by stimulating ARE-mediated TNF-alpha mRNA decay and several other inflammatory ARE- containing mRNAs in interferon (IFN)- and/or lipopolysaccharide (LPS)- induced macrophages (By similarity). Also plays a role in the regulation of dendritic cell maturation at the post-transcriptional level, and hence operates as part of a negative feedback loop to limit the inflammatory response (PubMed:[18367721](#)). Promotes ARE-mediated mRNA decay of hypoxia-inducible factor HIF1A mRNA during the response of endothelial cells to hypoxia (PubMed:[21775632](#)). Positively regulates early adipogenesis of preadipocytes by promoting ARE-mediated mRNA decay of immediate early genes (IEGs) (By similarity). Negatively regulates hematopoietic/erythroid cell differentiation by promoting ARE-mediated mRNA decay of the transcription factor STAT5B mRNA (PubMed:[20702587](#)). Plays a role in maintaining skeletal muscle satellite cell quiescence by promoting ARE-mediated mRNA decay of the myogenic determination factor MYOD1 mRNA (By similarity). Associates also with and regulates the expression of non-ARE-containing target mRNAs at the post-transcriptional level, such as MHC class I mRNAs (PubMed:[18367721](#)). Participates in association with argonaute RISC catalytic components in the ARE-mediated mRNA decay mechanism; assists microRNA (miRNA) targeting ARE-containing mRNAs (PubMed:[15766526](#)). May also play a role in the regulation of cytoplasmic mRNA decapping; enhances decapping of ARE-containing RNAs, in vitro (PubMed:[16364915](#)). Involved in the delivery of target ARE-mRNAs to processing bodies (PBs) (PubMed:[17369404](#)). In addition to its cytosolic mRNA-decay function, affects nuclear pre-mRNA processing (By similarity). Negatively regulates nuclear poly(A)-binding protein PABPN1-stimulated polyadenylation activity on ARE-containing pre-mRNA during LPS- stimulated macrophages (By similarity). Also involved in the regulation of stress granule (SG) and P-body (PB) formation and fusion (By similarity). Plays a role in the regulation of keratinocyte proliferation, differentiation and apoptosis (PubMed:[27182009](#)). Plays a role as a tumor suppressor by inhibiting cell proliferation in breast cancer cells (PubMed:[26926077](#)).

Cellular Location

Nucleus. Cytoplasm. Cytoplasmic granule. Cytoplasm, P-body. Note=Shuttles between nucleus and cytoplasm in a CRM1-dependent manner (By similarity). Localized predominantly in the cytoplasm in a p38 MAPK- and YWHAB-dependent manner (By similarity). Colocalizes with SH3KBP1 and MAP3K4 in the cytoplasm (PubMed:[20221403](#)). Component of cytoplasmic stress granules (SGs) (By similarity). Localizes to cytoplasmic stress granules upon energy starvation (PubMed:[15014438](#)). Localizes in processing bodies (PBs) (PubMed:[17369404](#)). Excluded from stress granules in a phosphorylation MAPKAPK2-dependent manner (By similarity). Shuttles in and out of both cytoplasmic P-body and SGs (By similarity) {ECO:0000250|UniProtKB:P22893, ECO:0000269|PubMed:[15014438](#), ECO:0000269|PubMed:[17369404](#), ECO:0000269|PubMed:[20221403](#)}

Tissue Location


Expressed in both basal and suprabasal epidermal layers (PubMed:[27182009](#)).

Expressed in epidermal keratinocytes (PubMed:27182009). Expressed strongly in mature dendritic cells (PubMed:18367721). Expressed in immature dendritic cells (at protein level) (PubMed:18367721).

References

Cao,H., et al., (2006) Biochem. J. 394 (PT 1), 285-297
Reconstitution and Storage:For short term use, store at 2-8C up to 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.

Images



WB Suggested Anti-ZFP36 Antibody Titration: 0.2-1 µg/ml
ELISA Titer: 1:312500
Positive Control: Human Lung

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.