

SMAD1 antibody - N-terminal region

Rabbit Polyclonal Antibody

Catalog # AI11444

Product Information

Application	WB, IHC
Primary Accession	Q99717
Other Accession	NM_005900 , NP_005891
Reactivity	Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Dog, Horse, Bovine
Predicted	Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Chicken, Dog, Horse, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	52258

Additional Information

Gene ID	4090
Alias Symbol	BSP1, JV4-1, JV41, MADH1, MADR1, BSP-1
Other Names	Mothers against decapentaplegic homolog 5, MAD homolog 5, Mothers against DPP homolog 5, JV5-1, SMAD family member 5, SMAD 5, Smad5, hSmad5, SMAD5, MADH5
Target/Specificity	This antibody reacts with SMAD1 + SMAD5 and to a lesser extent, SMAD8.
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 100 ul of distilled water. Final anti-SMAD1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
Precautions	SMAD1 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	SMAD5 (HGNC:6771)
Synonyms	MADH5
Function	Transcriptional regulator that plays a role in various cellular processes including embryonic development, cell differentiation, angiogenesis and tissue homeostasis (PubMed: 12064918 , PubMed: 16516194). Upon BMP ligand binding to their receptors at the cell surface, is phosphorylated by activated type I BMP receptors (BMPRIs) and associates with SMAD4 to form a

heteromeric complex which translocates into the nucleus acting as transcription factor (PubMed:[9442019](#)). In turn, the hetero-trimeric complex recognizes cis- regulatory elements containing Smad Binding Elements (SBEs) to modulate the outcome of the signaling network (PubMed:[33510867](#)). Non-phosphorylated SMAD5 has a cytoplasmic role in energy metabolism regulation by promoting mitochondrial respiration and glycolysis in response to cytoplasmic pH changes (PubMed:[28675158](#)). Mechanistically, interacts with hexokinase 1/HK1 and thereby accelerates glycolysis (PubMed:[28675158](#)).

Cellular Location

Cytoplasm. Nucleus Mitochondrion. Note=Cytoplasmic in the absence of ligand. Migrates to the nucleus when complexed with SMAD4

Tissue Location


Ubiquitous.

References

Jadlowiec,J.A., (2006) J. Biol. Chem. 281 (9), 5341-5347 Reconstitution and Storage: For short term use, store at 2-8C up to 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.

Images

Immunohistochemistry with Human Skin lysate tissue



SMAD1 antibody - N-terminal region (AI11444) validated by WB using Transfected 293T cell lysate at 2.5ug/ul.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.