

NAB2 antibody - N-terminal region

Rabbit Polyclonal Antibody

Catalog # AI11449

Product Information

Application	WB
Primary Accession	Q15742
Other Accession	NM_005967 , NP_005958
Reactivity	Human, Rat, Rabbit, Pig, Dog, Horse, Bovine
Predicted	Human, Rabbit, Pig, Horse, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	56594

Additional Information

Gene ID	4665
Alias Symbol	MADER
Other Names	NGFI-A-binding protein 2, EGR-1-binding protein 2, Melanoma-associated delayed early response protein, Protein MADER, NAB2, MADER
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 100 ul of distilled water. Final anti-NAB2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
Precautions	NAB2 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.


Protein Information

Name	NAB2
Synonyms	MADER
Function	Acts as a transcriptional repressor for zinc finger transcription factors EGR1 and EGR2. Isoform 2 lacks repression ability (By similarity).
Cellular Location	Nucleus. Note=Isoform 2 is not localized to the nucleus.
Tissue Location	Widely expressed at low levels. Highly expressed in melanoma cell lines

References

Srinivasan,R., et al., (er) J. Biol. Chem. (2006) In press
Reconstitution and Storage: For short term use, store at 2-8C up to 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.

Images



WB Suggested Anti-NAB2 Antibody Titration: 1.25µg/ml
ELISA Titer: 1:1562500
Positive Control: HepG2 cell lysate

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.