

PDLIM5 antibody - N-terminal region

Rabbit Polyclonal Antibody Catalog # AI11462

Product Information

Application	WB
Primary Accession	<u>Q96HC4</u>
Other Accession	<u>NM_006457, NP_006448</u>
Reactivity	Human, Mouse, Rat, Rabbit, Dog, Horse, Bovine
Predicted	Human, Mouse, Rat, Pig, Chicken, Dog, Horse, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	63945

Additional Information

Gene ID	10611
Alias Symbol Other Names	ENH, ENH1, L9, LIM PDZ and LIM domain protein 5, Enigma homolog, Enigma-like PDZ and LIM domains protein, PDLIM5, ENH
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 50 ul of distilled water. Final anti-PDLIM5 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
Precautions	PDLIM5 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	PDLIM5 {ECO:0000303 PubMed:15346770, ECO:0000312 HGNC:HGNC:17468}
Function	May play an important role in the heart development by scaffolding PKC to the Z-disk region. May play a role in the regulation of cardiomyocyte expansion. Isoforms lacking the LIM domains may negatively modulate the scaffolding activity of isoform 1. Overexpression promotes the development of heart hypertrophy. Contributes to the regulation of dendritic spine morphogenesis in neurons. May be required to restrain postsynaptic growth of excitatory synapses. Isoform 1, but not isoform 2, expression favors spine thinning and elongation.

Cellular Location	Postsynaptic density {ECO:0000250 UniProtKB:Q62920}. Presynapse {ECO:0000250 UniProtKB:Q62920}. Postsynapse {ECO:0000250 UniProtKB:Q62920}. Cytoplasm, cytosol {ECO:0000250 UniProtKB:Q62920}. Note=Detected both at presynaptic and postsynaptic sites, exclusively at excitatory synapses, but not inhibitory synapses, in hippocampal neurons {ECO:0000250 UniProtKB:Q62920}
Tissue Location	Heart and skeletal muscle specific. Expression is commonly increased in the brain of patients with bipolar disorder, schizophrenia, and major depression.

References

Squassina,A., (2008) Psychiatr. Genet. 18 (3), 128-132 Reconstitution and Storage:For short term use, store at 2-8C up to 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.

Images







Host: Rabbit Target Name: PDLIM5 Sample Tissue: Human Fetal Lung Antibody Dilution: 1.0µg/ml

PDLIM5



Host: Rabbit Target Name: PDLIM5 Sample Tissue: Human Fetal Liver Antibody Dilution: 1.0µg/ml

PDLIM5



Host: Rabbit Target Name: PDLIM5 Sample Tissue: Human Adult Placenta Antibody Dilution: 1.0µg/ml



Host: Rabbit Target Name:PDLIM5 Sample Tissue:HepG2 Lane A: Primary Antibody Lane B: Primary Antibody + Blocking Peptide Primary Antibody Concentration:1ug/ml Peptide Concentration: 5.0 ug/ml Lysate Quantity: 25ug/lane/lane Gel Concentration: 12%There is BioGPS gene expression data showing that PDLIM5 is expressed in HepG2

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.