

PDLIM5 antibody - N-terminal region

Rabbit Polyclonal Antibody Catalog # AI11462

Product Information

Application WB
Primary Accession Q96HC4

Other Accession NM 006457, NP 006448

ReactivityHuman, Mouse, Rat, Rabbit, Dog, Horse, Bovine **Predicted**Human, Mouse, Rat, Pig, Chicken, Dog, Horse, Bovine

Host Rabbit
Clonality Polyclonal
Calculated MW 63945

Additional Information

Gene ID 10611

Alias Symbol ENH, ENH1, L9, LIM

Other Names PDZ and LIM domain protein 5, Enigma homolog, Enigma-like PDZ and LIM

domains protein, PDLIM5, ENH

Format Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium

azide and 2% sucrose.

Reconstitution & Storage Add 50 ul of distilled water. Final anti-PDLIM5 antibody concentration is 1

mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at

20°C. Avoid repeat freeze-thaw cycles.

Precautions PDLIM5 antibody - N-terminal region is for research use only and not for use

in diagnostic or therapeutic procedures.

Protein Information

Name PDLIM5 {ECO:0000303 | PubMed:15346770,

ECO:0000312 | HGNC:HGNC:17468}

Function May play an important role in the heart development by scaffolding PKC to

the Z-disk region. May play a role in the regulation of cardiomyocyte expansion. Isoforms lacking the LIM domains may negatively modulate the scaffolding activity of isoform 1. Overexpression promotes the development

of heart hypertrophy. Contributes to the regulation of dendritic spine morphogenesis in neurons. May be required to restrain postsynaptic growth of excitatory synapses. Isoform 1, but not isoform 2, expression favors spine

thinning and elongation.

Cellular Location Postsynaptic density {ECO:0000250 | UniProtKB:Q62920}. Presynapse

{ECO:0000250|UniProtKB:Q62920}. Postsynapse {ECO:0000250|UniProtKB:Q62920}. Cytoplasm, cytosol

{ECO:0000250|UniProtKB:Q62920}. Note=Detected both at presynaptic and postsynaptic sites, exclusively at excitatory synapses, but not inhibitory synapses, in hippocampal neurons {ECO:0000250|UniProtKB:Q62920}

Tissue Location

Heart and skeletal muscle specific. Expression is commonly increased in the brain of patients with bipolar disorder, schizophrenia, and major depression.

References

Squassina, A., (2008) Psychiatr. Genet. 18 (3), 128-132 Reconstitution and Storage: For short term use, store at 2-8C up to 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.

Images

90 kDa_ 65 kDa_ 40 kDa_ 31 kDa_ 22 kDa_

WB Suggested Anti-PDLIM5 Antibody Titration: 0.2-1

μg/ml

ELISA Titer: 1:312500

Positive Control: HepG2 cell lysate

There is BioGPS gene expression data showing that

PDLIM5 is expressed in HepG2



Antibody Concentration: 1ug/mL

Host: Rabbit


Target Name: PDLIM5 Sample Tissue: HepG2

Antibody Dilution: 1.0µg/mlThere is BioGPS gene expression data showing that PDLIM5 is expressed in

HepG2

PDLIM5

22




Host: Rabbit

Target Name: PDLIM5

Sample Tissue: Human Fetal Brain Antibody Dilution: 1.0µg/ml


PDLIM5



Host: Rabbit

Target Name: PDLIM5


Sample Tissue: Human Fetal Heart Antibody Dilution: 1.0µg/ml



Host: Rabbit

Target Name: PDLIM5


Sample Tissue: Human Fetal Lung Antibody Dilution: 1.0µg/ml



Host: Rabbit

Target Name: PDLIM5

Sample Tissue: Human Fetal Liver Antibody Dilution: 1.0µg/ml




Host: Rabbit

Target Name: PDLIM5

Sample Tissue: Human Adult Placenta

Antibody Dilution: 1.0µg/ml



Host: Rabbit Target Name:PDLIM5 Sample Tissue:HepG2 Lane A: Primary Antibody Lane B: Primary Antibody + Blocking Peptide Primary Antibody Concentration:1ug/ml Peptide Concentration: 5.0 ug/ml Lysate Quantity: 25ug/lane/lane Gel Concentration: 12%There is BioGPS gene expression data showing that PDLIM5 is expressed in HepG2

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.