

HMG20A antibody - C-terminal region

Rabbit Polyclonal Antibody

Catalog # AI11534

Product Information

Application	WB
Primary Accession	Q9NP66
Other Accession	NM_018200 , NP_060670
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Dog, Horse, Bovine
Predicted	Human, Rat, Rabbit, Pig, Chicken, Dog, Horse, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	40144

Additional Information

Gene ID	10363
Alias Symbol	HMGX1, HMGXB1
Other Names	High mobility group protein 20A, HMG box-containing protein 20A, HMG domain-containing protein 1, HMG domain-containing protein HMGX1, HMG20A, HMGX1, HMGXB1
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 50 ul of distilled water. Final anti-HMG20A antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
Precautions	HMG20A antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	HMG20A
Synonyms	HMGX1, HMGXB1
Function	Plays a role in neuronal differentiation as chromatin- associated protein. Acts as inhibitor of HMG20B. Overcomes the repressive effects of the neuronal silencer REST and induces the activation of neuronal-specific genes. Involved in the recruitment of the histone methyltransferase KMT2A/MLL1 and consequent increased methylation of histone H3 lysine 4 (By similarity).
Cellular Location	Nucleus {ECO:0000255 PROSITE-ProRule:PRU00267}.


Tissue Location

Ubiquitous..

References

Rual,J.F., et al., (2005) Nature 437 (7062), 1173-1178
Reconstitution and Storage:For short term use, store at 2-8C up to 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.

Images



WB Suggested Anti-HMG20A Antibody Titration: 0.2-1
µg/ml
ELISA Titer: 1:1562500
Positive Control: Jurkat cell lysate

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.