

TRIM62 antibody - N-terminal region

Rabbit Polyclonal Antibody Catalog # AI11535

Product Information

Application WB
Primary Accession Q9BVG3

Other Accession <u>NM_018207</u>, <u>NP_060677</u>

Reactivity Human, Mouse, Rat, Pig, Horse, Bovine **Predicted** Human, Mouse, Dog, Horse, Bovine

Host Rabbit
Clonality Polyclonal
Calculated MW 54268

Additional Information

Gene ID 55223

Alias Symbol DEAR1

Other Names E3 ubiquitin-protein ligase TRIM62, 6.3.2.-, Tripartite motif-containing protein

62, TRIM62

Format Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium

azide and 2% sucrose.

Reconstitution & Storage Add 100 ul of distilled water. Final anti-TRIM62 antibody concentration is 1

mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at

20°C. Avoid repeat freeze-thaw cycles.

Precautions TRIM62 antibody - N-terminal region is for research use only and not for use

in diagnostic or therapeutic procedures.

Protein Information

Name TRIM62 {ECO:0000303 | PubMed:23402750,

ECO:0000312 | HGNC:HGNC:25574}

Function E3 ubiquitin ligase that plays a role in antifungal immunity by mediating

'Lys-27'-linked ubiquitination of CARD9 downstream of C- type lectin

receptors; leading to CARD9 activation, followed by activation of NF-kappa-B and MAP kinase p38 pathways (PubMed: 26488816). E3 ubiquitin ligase activity

is dependent on E2 ubiquitin-conjugating enzyme UBE2D2


(PubMed: 23402750).

Cellular Location Cytoplasm

References

Isogai, T., et al., (2000) Unpublished Reconstitution and Storage: For short term use, store at 2-8C up to 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.

Images



WB Suggested Anti-TRIM62 Antibody Titration: 1.25µg/ml

ELISA Titer: 1:1562500

Positive Control: Jurkat cell lysate

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.