

Klhl12 antibody - N-terminal region

Rabbit Polyclonal Antibody

Catalog # AI11564

Product Information

Application	WB
Primary Accession	Q8R2H4
Other Accession	NM_153730 , NP_714952
Reactivity	Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Dog, Horse, Bovine
Predicted	Human, Rat, Rabbit, Zebrafish, Pig, Dog, Horse, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	63306

Additional Information

Gene ID	266772
Alias Symbol	C3ip1, MGC93127
Other Names	Kelch-like protein 12, CUL3-interacting protein 1, Klhl12, C3ip1
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 50 ul of distilled water. Final anti-Klhl12 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
Precautions	Klhl12 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information


Name	Klhl12
Synonyms	C3ip1
Function	Substrate-specific adapter of a BCR (BTB-CUL3-RBX1) E3 ubiquitin ligase complex that acts as a negative regulator of Wnt signaling pathway and ER-Golgi transport. The BCR(KLHL12) complex is involved in ER-Golgi transport by regulating the size of COPII coats, thereby playing a key role in collagen export, which is required for embryonic stem (ES) cells division: BCR(KLHL12) acts by mediating monoubiquitination of SEC31 (SEC31A or SEC31B). The BCR(KLHL12) complex is also involved in neural crest specification: in response to cytosolic calcium increase, interacts with the heterodimer formed with PEF1 and PDCD6/ALG-2, leading to bridge together the BCR(KLHL12)

complex and SEC31 (SEC31A or SEC31B), promoting monoubiquitination of SEC31 and subsequent collagen export. As part of the BCR(KLHL12) complex, also acts as a negative regulator of the Wnt signaling pathway by mediating ubiquitination and subsequent proteolysis of DVL3. The BCR(KLHL12) complex also mediates polyubiquitination of DRD4 and PEF1, without leading to degradation of these proteins.

Cellular Location

Cytoplasmic vesicle, COPII-coated vesicle {ECO:0000250|UniProtKB:Q53G59}

Images



WB Suggested Anti-Klhl12 Antibody Titration: 1.0 µg/ml
Positive Control: Rat Brain

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.