

SIRT5 antibody - C-terminal region

Rabbit Polyclonal Antibody Catalog # AI11677

Product Information

Application WB Primary Accession Q5T295

Other Accession <u>NM 006800, NP 112534</u>

Reactivity Human, Mouse, Rat, Rabbit, Zebrafish, Goat, Dog, Horse, Bovine

Predicted Human, Mouse, Pig, Chicken, Goat, Bovine

Host Rabbit
Clonality Polyclonal
Calculated MW 33 KDa

Additional Information

Alias Symbol SIR2L5

Format Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium

azide and 2% sucrose.

Reconstitution & Storage Add 100 ul of distilled water. Final anti-SIRT5 antibody concentration is 1

mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at

20°C. Avoid repeat freeze-thaw cycles.

Precautions SIRT5 antibody - C-terminal region is for research use only and not for use in

diagnostic or therapeutic procedures.

Protein Information

References


Mahlknecht, U., et al., (2006) Cytogenet. Genome Res. 112 (3-4), 208-212Reconstitution and Storage: For short term use, store at 2-8C up to 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.


Images

WB Suggested Anti-SIRT5 Antibody Titration: 5.0µg/ml

ELISA Titer: 1:62500

Positive Control: Human heart





See Immunoblot 2 Data and Customer Feedback for more information

Lanes: Lane1: Untransfected HEK293T lysate Lane2:

mSIRT5-FLAG transfected HEK293T lysate

Primary Antibody Dilution: 1:200 Secondary Antibody: ProteinA-HRP Secondary Antibody Dilution: 1:5000

Gene Name: SIRT5

Submitted by: Anonymous There is BioGPS gene expression data showing that SIRT5 is expressed in

HEK293T

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.