

# DDX19B antibody - C-terminal region

Rabbit Polyclonal Antibody Catalog # AI11723

#### **Product Information**

Application WB
Primary Accession Q9UMR2

Other Accession <u>NM\_007242</u>, <u>NP\_009173</u>

**Reactivity** Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Dog, Horse, Bovine, Yeast

**Predicted** Human, Mouse, Rat, Rabbit, Pig, Chicken, Bovine

Host Rabbit
Clonality Polyclonal
Calculated MW 53927

### **Additional Information**

**Gene ID** 11269

Alias Symbol DBP5, RNAh, DDX19

Other Names ATP-dependent RNA helicase DDX19B, 3.6.4.13, DEAD box RNA helicase

DEAD5, DEAD box protein 19B, DDX19B, DBP5, DDX19, TDBP

Format Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium

azide and 2% sucrose.

**Reconstitution & Storage** Add 100 ul of distilled water. Final anti-DDX19B antibody concentration is 1

mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at

20°C. Avoid repeat freeze-thaw cycles.

**Precautions** DDX19B antibody - C-terminal region is for research use only and not for use

in diagnostic or therapeutic procedures.

#### **Protein Information**

Name DDX19B

**Synonyms** DBP5, DDX19, TDBP

**Function** ATP-dependent RNA helicase involved in mRNA export from the nucleus

(PubMed: 10428971). Rather than unwinding RNA duplexes, DDX19B functions as a remodeler of ribonucleoprotein particles, whereby proteins bound to nuclear mRNA are dissociated and replaced by cytoplasmic mRNA binding

proteins (PubMed: 10428971).


**Cellular Location** Cytoplasm. Nucleus, nucleoplasm. Note=Associates with the nuclear pore

complex cytoplasmic fibrils

## References

Yin,L., Reprod. Fertil. Dev. 14 (3-4), 185-189 (2002)Reconstitution and Storage:For short term use, store at 2-8C up to 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.

# **Images**



WB Suggested Anti-DDX19B Antibody Titration: 2.5µg/ml Positive Control: HepG2 cell lysate DDX19B is supported by BioGPS gene expression data to be expressed in HepG2

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.