

Cstf2 antibody - N-terminal region

Rabbit Polyclonal Antibody

Catalog # AI11735

Product Information

Application	WB
Primary Accession	Q8BIQ5
Other Accession	NM_133196 , NP_573459
Reactivity	Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Dog, Horse, Bovine
Predicted	Human, Mouse, Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	61341

Additional Information

Gene ID	108062
Alias Symbol	64kDa, C630034J23Rik, Cstf64
Other Names	Cleavage stimulation factor subunit 2, CF-1 64 kDa subunit, Cleavage stimulation factor 64 kDa subunit, CSTF 64 kDa subunit, CstF-64, Cstf2
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 50 ul of distilled water. Final anti-Cstf2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
Precautions	Cstf2 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	Cstf2
Function	One of the multiple factors required for polyadenylation and 3'-end cleavage of mammalian pre-mRNAs. This subunit is directly involved in the binding to pre-mRNAs (By similarity).
Cellular Location	Nucleus. Note=Localized with DDX1 in cleavage bodies {ECO:0000250 UniProtKB:P33240}
Tissue Location	Expressed in most somatic cell types (at protein level). Highly expressed in testis, except in meiotic spermatocytes

Images



WB Suggested Anti-Cstf2 Antibody Titration: 1.0 µg/ml
Positive Control: Mouse Pancreas

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.