

# HNRPA1 antibody - N-terminal region

Rabbit Polyclonal Antibody Catalog # AI11739

#### **Product Information**

Application WB, IHC Primary Accession Q6IPF2

Other Accession NM 002136, NP 002127

**Reactivity**Human, Mouse, Rat, Rabbit, Pig, Dog, Horse, Bovine **Predicted**Human, Mouse, Rat, Pig, Chicken, Dog, Horse, Bovine

Host Rabbit
Clonality Polyclonal
Calculated MW 35 KDa

## **Additional Information**

Alias Symbol HNRPA1, HNRPA1L3, hnRNP A1, hnRNP-A1

Format Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium

azide and 2% sucrose.

**Reconstitution & Storage** Add 100 ul of distilled water. Final anti-HNRPA1 antibody concentration is 1

mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at

20°C. Avoid repeat freeze-thaw cycles.

**Precautions** HNRPA1 antibody - N-terminal region is for research use only and not for use


in diagnostic or therapeutic procedures.

### **Protein Information**

#### References

Tevosian,S.G., (2005) J. Biol. Chem. 280 (13), 13171-13178Reconstitution and Storage:For short term use, store at 2-8C up to 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.Publications:Li, S. et al. Identification of an aptamer targeting hnRNP A1 by tissue slide-based SELEX. J. Pathol. 218, 327-36 (2009). ICC/IF, H, Human, Rat, Dog, Pig, Rabbit, Guinea pig, Mouse, Bovine19291713Gonen, N. & Assaraf, Y. G. The obligatory intestinal folate transporter PCFT (SLC46A1) is regulated by nuclear respiratory factor 1. J. Biol. Chem. 285, 33602-13 (2010). WB, H, Human, Rat, Dog, Pig, Rabbit, Guinea pig, Mouse, Bovine20724482Stark, M., Bram, E. E., Akerman, M., Mandel-Gutfreund, Y. & Assaraf, Y. G. Heterogeneous nuclear ribonucleoprotein H1/H2-dependent unsplicing of thymidine phosphorylase results in anticancer drug resistance. J. Biol. Chem. 286, 3741-54 (2011). WB, ICC/IF, ChIP, H, Human, Rat, Dog, Pig, Rabbit, Guinea pig, Mouse, Bovine21068389Li, S. et al. Aptamer BC15 against heterogeneous nuclear ribonucleoprotein A1 has potential value in diagnosis and therapy of hepatocarcinoma. Nucleic Acid Ther. 22, 391-8 (2012). WB, IHC, H, Human, Rat, Dog, Pig, Rabbit, Guinea pig, Mouse, Bovine23062008

## **Images**




Rabbit Anti-HNRPA1 Antibody
Paraffin Embedded Tissue: Human Liver

Cellular Data: Hepatocytes

Antibody Concentration: 4.0-8.0 μg/ml

Magnification: 400X



Rabbit Anti-HNRPA1 Antibody Paraffin Embedded Tissue: Human Heart Cellular Data: Myocardial cells


Antibody Concentration: 4.0-8.0 μg/ml

Magnification: 400X



Rabbit Anti-HNRPA1 antibody Paraffin Embedded Tissue: Human Heart cell Cellular Data: cardiac cell of renal tubule Antibody Concentration: 4.0-8.0 µg/ml

Magnification: 400X



WB Suggested Anti-HNRPA1 Antibody Titration: 1.25µg/ml

ELISA Titer: 1:312500

Positive Control: Jurkat cell lysate

HNRNPA1 is strongly supported by BioGPS gene

expression data to be expressed in Human Jurkat cells

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.