

# PSMA1 antibody - N-terminal region

Rabbit Polyclonal Antibody

Catalog # AI11747

## Product Information

---

<b>Application</b>	WB
<b>Primary Accession</b>	<a href="#">P25786</a>
<b>Other Accession</b>	<a href="#">NM_002786</a> , <a href="#">NP_002777</a>
<b>Reactivity</b>	Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Dog, Horse, Bovine, Yeast
<b>Predicted</b>	Human, Mouse, Zebrafish, Chicken, Dog, Horse
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Calculated MW</b>	29556

## Additional Information

---

<b>Gene ID</b>	5682
<b>Alias Symbol</b>	NU, HC2, PROS30
<b>Other Names</b>	Proteasome subunit alpha type-1, 3.4.25.1, 30 kDa prosomal protein, PROS-30, Macropain subunit C2, Multicatalytic endopeptidase complex subunit C2, Proteasome component C2, Proteasome nu chain, PSMA1, HC2, NU, PROS30, PSC2
<b>Format</b>	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
<b>Reconstitution &amp; Storage</b>	Add 100 ul of distilled water. Final anti-PSMA1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
<b>Precautions</b>	PSMA1 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	PSMA1 ( <a href="#">HGNC:9530</a> )
<b>Synonyms</b>	HC2, NU, PROS30, PSC2
<b>Function</b>	Component of the 20S core proteasome complex involved in the proteolytic degradation of most intracellular proteins. This complex plays numerous essential roles within the cell by associating with different regulatory particles. Associated with two 19S regulatory particles, forms the 26S proteasome and thus participates in the ATP- dependent degradation of ubiquitinated proteins. The 26S proteasome plays a key role in the

maintenance of protein homeostasis by removing misfolded or damaged proteins that could impair cellular functions, and by removing proteins whose functions are no longer required. Associated with the PA200 or PA28, the 20S proteasome mediates ubiquitin- independent protein degradation. This type of proteolysis is required in several pathways including spermatogenesis (20S-PA200 complex) or generation of a subset of MHC class I-presented antigenic peptides (20S-PA28 complex).

#### Cellular Location

Cytoplasm. Nucleus. Note=Translocated from the cytoplasm into the nucleus following interaction with AKIRIN2, which bridges the proteasome with the nuclear import receptor IPO9


## References

---

Conticello,S.G., (2003) Curr. Biol. 13 (22), 2009-2013  
Reconstitution and Storage:For short term use, store at 2-8C up to 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.

## Images

---



WB Suggested Anti-PSMA1 Antibody Titration: 1.25µg/ml  
Positive Control: Jurkat cell lysate

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.