

SLC12A1 antibody - N-terminal region

Rabbit Polyclonal Antibody

Catalog # AI11886

Product Information


Application	WB, IHC
Primary Accession	Q8IUN5
Other Accession	NM_000338 , AAH40138
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Dog, Horse, Bovine
Predicted	Human, Mouse, Rat, Pig, Chicken, Dog, Horse, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	47 KDa

Additional Information

Alias Symbol	BSC1, MGC48843, NKCC2
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 50 ul of distilled water. Final anti-SLC12A1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
Precautions	SLC12A1 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.


Protein Information

Images




Rabbit Anti-SLC12A1 Antibody
Paraffin Embedded Tissue: Human Kidney
Cellular Data: Epithelial cells of renal tubule
Antibody Concentration: 4.0-8.0 µg/ml
Magnification: 400X

WB Suggested Anti-SLC12A1 Antibody Titration: 0.2-1 µg/ml



Positive Control: Jurkat cell lysate



Host: Rabbit
Target Name: SLC12A1
Sample Tissue: Jurkat
Lane A: Primary Antibody
Lane B: Primary Antibody + Blocking Peptide
Primary Antibody
Concentration: 1 µg/ml
Peptide Concentration: 5 µg/ml
Lysate Quantity: 25 µg/lane/lane Gel
Concentration: 0.12

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.