

TSHR antibody - C-terminal region

Rabbit Polyclonal Antibody

Catalog # AI11952

Product Information

Application	WB
Primary Accession	Q96GT6
Other Accession	NM_001018036 , NP_001018046
Reactivity	Human, Mouse, Rat, Rabbit, Dog, Horse, Bovine, Sheep
Predicted	Human, Rabbit, Pig, Dog, Horse, Bovine, Sheep
Host	Rabbit
Clonality	Polyclonal
Calculated MW	28 KDa

Additional Information

Alias Symbol	LGR3, MGC75129, hTSHR-I, CHNG1
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 50 ul of distilled water. Final anti-TSHR antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
Precautions	TSHR antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.


Protein Information

References

Claus, M., (2005) Endocrinology 146 (12), 5197-5203
Reconstitution and Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.
Publications: Pessina, P., Castillo, V., Sartore, I., Borrego, J. & Meikle, A. Semiquantitative immunohistochemical marker staining and localization in canine thyroid carcinoma and normal thyroid gland. Vet. Comp. Oncol. (2014). doi:10.1111/vco.12111
IHC, Human, H, Rat, Dog, Rabbit, Guinea pig, Mouse, Bovine, Sheep 25082554

Images

WB Suggested Anti-TSHR Antibody Titration: 0.2-1 µg/ml
Positive Control: Jurkat cell lysate



There is BioGPS gene expression data showing that TSHR is expressed in Jurkat

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.