

FADS1 antibody - N-terminal region

Rabbit Polyclonal Antibody

Catalog # AI12012

Product Information

Application	WB
Primary Accession	O60427
Other Accession	NM_013402 , NP_037534
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Dog, Horse, Bovine
Predicted	Human, Mouse, Rat, Pig, Chicken, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	51964

Additional Information

Gene ID	3992
Alias Symbol	D5D, FADS6, FADSD5, FLJ38956, FLJ90273, LLCDL1, TU12
Other Names	Fatty acid desaturase 1, 1.14.19.-, Delta(5) fatty acid desaturase, D5D, Delta(5) desaturase, Delta-5 desaturase, FADS1, FADSD5
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 50 ul of distilled water. Final anti-FADS1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
Precautions	FADS1 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	FADS1 {ECO:0000303 PubMed:10860662, ECO:0000312 HGNC:HGNC:3574}
Function	[Isoform 1]: Acts as a front-end fatty acyl-coenzyme A (CoA) desaturase that introduces a cis double bond at carbon 5 located between a preexisting double bond and the carboxyl end of the fatty acyl chain. Involved in biosynthesis of highly unsaturated fatty acids (HUFA) from the essential polyunsaturated fatty acids (PUFA) linoleic acid (LA) (18:2n-6) and alpha-linolenic acid (ALA) (18:3n-3) precursors. Specifically, desaturates dihomo-gamma-linoleoate (DGLA) (20:3n-6) and eicosatetraenoate (ETA) (20:4n-3) to generate arachidonate (AA) (20:4n-6) and eicosapentaenoate (EPA) (20:5n-3), respectively (PubMed: 10601301 , PubMed: 10769175). As a rate limiting enzyme for DGLA (20:3n-6) and AA (20:4n-6)-derived eicosanoid

biosynthesis, controls the metabolism of inflammatory lipids like prostaglandin E2, critical for efficient acute inflammatory response and maintenance of epithelium homeostasis. Contributes to membrane phospholipid biosynthesis by providing AA (20:4n-6) as a major acyl chain esterified into phospholipids. In particular, regulates phosphatidylinositol-4,5-bisphosphate levels, modulating inflammatory cytokine production in T-cells (By similarity). Also desaturates (11E)-octadecenoate (trans-vaccenoate)(18:1n-9), a metabolite in the biohydrogenation pathway of LA (18:2n-6) (By similarity).

Cellular Location

[Isoform 1]: Endoplasmic reticulum membrane
{ECO:0000250|UniProtKB:A4UVI1}; Multi-pass membrane protein
{ECO:0000250|UniProtKB:A4UVI1}. Mitochondrion


Tissue Location

Widely expressed, with highest levels in liver, brain, adrenal gland and heart. Highly expressed in fetal liver and brain.

References

Rzehak,P., (er) Br. J. Nutr., 1-7 (2008) In press
Reconstitution and Storage:For short term use, store at 2-8C up to 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.

Images



WB Suggested Anti-FADS1 Antibody Titration: 0.2-1 µg/ml
ELISA Titer: 1:312500
Positive Control: 721_B cell lysate
FADS1 is strongly supported by BioGPS gene expression data to be expressed in Human 721_B cells

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.