

Dkk4 antibody - C-terminal region

Rabbit Polyclonal Antibody Catalog # AI12014

Product Information


Application	WB
Primary Accession	<u>Q8VEJ3</u>
Other Accession	<u>NM_145592</u> , <u>NP_663567</u>
Reactivity	Human, Mouse, Rat, Rabbit, Dog, Horse
Predicted	Human, Rat, Rabbit, Pig, Horse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	24261

Additional Information

Gene ID	234130
Alias Symbol Other Names	Dkk-4, MGC25705 Dickkopf-related protein 4, Dickkopf-4, Dkk-4, Dkk4
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 50 ul of distilled water. Final anti-Dkk4 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
Precautions	Dkk4 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	Dkk4
Function	Antagonizes canonical Wnt signaling by inhibiting LRP5/6 interaction with Wnt and by forming a ternary complex with the transmembrane protein KREMEN that promotes internalization of LRP5/6. DKKs play an important role in vertebrate development, where they locally inhibit Wnt regulated processes such as antero-posterior axial patterning, limb development, somitogenesis and eye formation. In the adult, Dkks are implicated in bone formation and bone disease, cancer and Alzheimer disease (By similarity).
Cellular Location	Secreted.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.