

JUND antibody - C-terminal region

Rabbit Polyclonal Antibody

Catalog # AI12074

Product Information

Application	WB
Primary Accession	P17535
Other Accession	NM_005354 , NP_005345
Reactivity	Human, Mouse, Rat, Rabbit, Zebrafish, Goat, Dog, Guinea Pig, Horse, Bovine, Sheep
Predicted Host	Human, Mouse, Rat, Zebrafish, Chicken, Dog, Guinea Pig, Bovine
Clonality	Rabbit
Calculated MW	Polyclonal 35174

Additional Information

Gene ID	3727
Alias Symbol	AP-1
Other Names	Transcription factor jun-D, JUND
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 50 ul of distilled water. Final anti-JUND antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
Precautions	JUND antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.


Protein Information

Name	JUND
Function	Transcription factor binding AP-1 sites (PubMed: 9989505). Heterodimerizes with proteins of the FOS family to form an AP-1 transcription factor complex, thereby enhancing their DNA binding activity to an AP-1 consensus sequence 3'-TGA[GC]TCA-5' and enhancing their transcriptional activity (PubMed: 28981703 , PubMed: 9989505).
Cellular Location	Nucleus.

References

Ota,T.,etal.,(2005)Oncogene24(51),7567-7578ReconstitutionandStorage:Forshorttermuse,storeat2-8Cupto1week.Forlongtermstorage,storeat-20Cinsmallaliquotstopreventfreeze-thawcycles.

Images



WB Suggested Anti-JUND Antibody Titration: 0.2-1 µg/ml
ELISA Titer: 1:62500
Positive Control: Human Placenta

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.