

# UTX antibody - N-terminal region

Rabbit Polyclonal Antibody Catalog # AI12082

#### **Product Information**

Application WB Primary Accession 015550

Other Accession <u>NM 021140, NP 066963</u>

**Reactivity** Human, Mouse, Rat, Zebrafish, Pig, Dog, Bovine

**Predicted** Human, Zebrafish, Pig

Host Rabbit
Clonality Polyclonal
Calculated MW 154177

### **Additional Information**

**Gene ID** 7403

**Alias Symbol** DKFZp686A03225, MGC141941, UTX, bA386N14.2, KABUK2

Other Names Lysine-specific demethylase 6A, 1.14.11.-, Histone demethylase UTX,

Ubiquitously-transcribed TPR protein on the X chromosome,

Ubiquitously-transcribed X chromosome tetratricopeptide repeat protein,

KDM6A, UTX

Format Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium

azide and 2% sucrose.

**Reconstitution & Storage** Add 50 ul of distilled water. Final anti-UTX antibody concentration is 1 mg/ml

in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C.

Avoid repeat freeze-thaw cycles.

**Precautions**UTX antibody - N-terminal region is for research use only and not for use in

diagnostic or therapeutic procedures.

## **Protein Information**

Name KDM6A

**Synonyms** UTX

**Function** Histone demethylase that specifically demethylates 'Lys-27' of histone H3,

thereby playing a central role in histone code (PubMed: 17713478,

PubMed: 17761849, PubMed: 17851529). Demethylates trimethylated and dimethylated but not monomethylated H3 'Lys-27' (PubMed: 17713478, PubMed: 17761849, PubMed: 17851529). Plays a central role in regulation of


posterior development, by regulating HOX gene expression

(PubMed: <u>17851529</u>). Demethylation of 'Lys-27' of histone H3 is concomitant with methylation of 'Lys-4' of histone H3, and regulates the recruitment of the PRC1 complex and monoubiquitination of histone H2A (PubMed: <u>17761849</u>). Plays a demethylase-independent role in chromatin remodeling to regulate T-box family member-dependent gene expression (By similarity).

#### **Cellular Location**

Nucleus.

# **Images**



WB Suggested Anti-UTX Antibody Titration: 0.2-1 µg/ml

ELISA Titer: 1:1562500

Positive Control: Hela cell lysate

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.