

# RAB38 antibody - N-terminal region

Rabbit Polyclonal Antibody

Catalog # AI12126

## Product Information

<b>Application</b>	WB
<b>Primary Accession</b>	<a href="#">P57729</a>
<b>Other Accession</b>	<a href="#">NM_022337</a> , <a href="#">NP_071732</a>
<b>Reactivity</b>	Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Dog, Guinea Pig, Horse, Bovine
<b>Predicted</b>	Human, Mouse, Rat, Chicken, Dog
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Calculated MW</b>	23712

## Additional Information

<b>Gene ID</b>	23682
<b>Alias Symbol</b>	NY-MEL-1, rrGTPbp
<b>Other Names</b>	Ras-related protein Rab-38, Melanoma antigen NY-MEL-1, RAB38
<b>Format</b>	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
<b>Reconstitution &amp; Storage</b>	Add 50 ul of distilled water. Final anti-RAB38 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
<b>Precautions</b>	RAB38 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

<b>Name</b>	RAB38 ( <a href="#">HGNC:9776</a> )
<b>Function</b>	The small GTPases Rab are key regulators of intracellular membrane trafficking, from the formation of transport vesicles to their fusion with membranes. Rabs cycle between an inactive GDP-bound form and an active GTP-bound form that is able to recruit to membranes different sets of downstream effectors directly responsible for vesicle formation, movement, tethering and fusion (By similarity). RAB38 may be involved in melanosomal transport and docking. Involved in the proper sorting of TYRP1. Involved in peripheral melanosomal distribution of TYRP1 in melanocytes; the function, which probably is implicating vesicle-trafficking, includes cooperation with ANKRD27 and VAMP7 (By similarity). Plays a role in the maturation of phagosomes that engulf pathogens, such as S.aureus and M.tuberculosis

(PubMed:[21255211](#)). Plays an important role in the control of melanin production and melanosome biogenesis (PubMed:[23084991](#)). In concert with RAB32, regulates the proper trafficking of melanogenic enzymes TYR, TYRP1 and DCT/TYRP2 to melanosomes in melanocytes (By similarity).

## Cellular Location

Cell membrane; Lipid-anchor; Cytoplasmic side. Melanosome. Cytoplasmic vesicle, phagosome. Cytoplasmic vesicle, phagosome membrane; Lipid-anchor; Cytoplasmic side. Melanosome membrane Note=Recruited to phagosomes containing S.aureus or M.tuberculosis (PubMed:21255211). The BLOC-3 complex, a heterodimer of HPS1 and HPS4 promotes its membrane localization (PubMed:23084991)


## Tissue Location

Expressed in melanocytes.


## References

Wang, F., (2008) Biochem. Biophys. Res. Commun. 372(1), 162-167 Reconstitution and Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.

## Images



WB Suggested Anti-RAB38 Antibody Titration: 0.2-1 µg/ml  
ELISA Titer: 1:312500  
Positive Control: MCF7 cell lysate



Human , Mouse

See Immunoblot 2 Data and Customer Feedback for more information

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.