

Dcaf11 antibody - C-terminal region

Rabbit Polyclonal Antibody

Catalog # AI12135

Product Information

Application	WB
Primary Accession	Q9EQQ2
Other Accession	NM_023311 , NP_075800
Reactivity	Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Goat, Dog, Guinea Pig, Horse, Bovine
Predicted Host	Human, Mouse, Rabbit, Zebrafish, Dog, Bovine
Clonality	Rabbit
Calculated MW	Polyclonal 27873

Additional Information

Gene ID	67180
Alias Symbol Other Names	0710008A13Rik, C76035, D14Ucla1, Dcaf11, GLO14, Wdr23 Protein YIPF5, YIP1 family member 5, YPT-interacting protein 1 A, Yipf5, Yip1a
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 50 ul of distilled water. Final anti-Dcaf11 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
Precautions	Dcaf11 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information


Name	Yipf5 {ECO:0000312 MGI:MGI:1914430}
Synonyms	Yip1a
Function	Plays a role in transport between endoplasmic reticulum and Golgi. In pancreatic beta cells, required to transport proinsulin from endoplasmic reticulum into the Golgi.
Cellular Location	Golgi apparatus, cis-Golgi network membrane; Multi-pass membrane protein. Cytoplasmic vesicle, COPII-coated vesicle {ECO:0000250 UniProtKB:Q5XID0}. Endoplasmic reticulum membrane; Multi-pass membrane protein. Note=Enriched at the endoplasmic reticulum exit sites (PubMed:15254263).

Incorporated into COPII coated vesicles (By similarity).
{ECO:0000250|UniProtKB:Q5XID0, ECO:0000269|PubMed:15254263}

Tissue Location

Ubiquitously expressed.

Images



WB Suggested Anti-Dcaf11 Antibody Titration: 1.0 µg/ml
Positive Control: Mouse pancreas

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.