10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

# Rfwd2 antibody - middle region

Rabbit Polyclonal Antibody
Catalog # AI12193

## Product Information

| | |
|---|---|
| **Application** | WB |
| **Primary Accession** | Q9R1A8 |
| **Other Accession** | NM_011931, NP_036061 |
| **Reactivity** | Human, Mouse, Rat, Rabbit, Pig, Dog, Guinea Pig, Horse, Bovine |
| **Predicted** | Human, Mouse, Rat, Rabbit, Pig, Chicken, Dog, Guinea Pig, Horse, Bovine |
| **Host** | Rabbit |
| **Clonality** | Polyclonal |
| **Calculated MW** | 80441 |

## Additional Information

| | |
|---|---|
| **Gene ID** | 26374 |
| **Alias Symbol** | AI316802, C80879, Cop1 |
| **Other Names** | E3 ubiquitin-protein ligase RFWD2, 6.3.2.-, Constitutive photomorphogenesis protein 1 homolog, mCOP1, RING finger and WD repeat domain protein 2, Rfwd2, Cop1 |
| **Format** | Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose. |
| **Reconstitution & Storage** | Add 50 ul of distilled water. Final anti-Rfwd2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles. |
| **Precautions** | Rfwd2 antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures. |


## Protein Information

| | |
|---|---|
| **Name** | Cop1 {ECO:0000312\|MGI:MGI:1347046} |
| **Function** | E3 ubiquitin-protein ligase that mediates ubiquitination and subsequent proteasomal degradation of target proteins. E3 ubiquitin ligases accept ubiquitin from an E2 ubiquitin-conjugating enzyme in the form of a thioester and then directly transfers the ubiquitin to targeted substrates. Involved in JUN ubiquitination and degradation. Directly involved in p53 (TP53) ubiquitination and degradation, thereby abolishing p53-dependent transcription and apoptosis. Ubiquitinates p53 independently of MDM2 or RCHY1. Probably mediates E3 ubiquitin ligase activity by functioning as the essential RING domain subunit of larger E3 complexes. In contrast, it does not |

constitute the catalytic RING subunit in the DCX DET1-COP1 complex that negatively regulates JUN, the ubiquitin ligase activity being mediated by RBX1. Involved in 14-3-3 protein sigma/SFN ubiquitination and proteasomal degradation, leading to AKT activation and promotion of cell survival. Ubiquitinates MTA1 leading to its proteasomal degradation. Upon binding to TRIB1, ubiquitinates CEBPA, which lacks a canonical COP1-binding motif.

**Cellular Location** Nucleus speckle. Cytoplasm. Note=In the nucleus, it forms nuclear speckles

## Images




WB Suggested Anti-Rfwd2 Antibody Titration: 1.0 µg/ml
Positive Control: Mouse Thymus

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.