

MARCH2 Antibody - C-terminal region

Rabbit Polyclonal Antibody

Catalog # AI12195

Product Information

Application	WB
Primary Accession	Q9P0N8
Other Accession	NM_016496 , NP_057580
Reactivity	Human, Rabbit, Guinea Pig
Predicted	Human, Rabbit
Host	Rabbit
Clonality	Polyclonal
Calculated MW	26995

Additional Information

Gene ID	51257
Alias Symbol	MARCH-II, RNF172
Other Names	E3 ubiquitin-protein ligase MARCH2, 6.3.2.-, Membrane-associated RING finger protein 2, Membrane-associated RING-CH protein II, MARCH-II, RING finger protein 172, MARCH2, RNF172
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 50 μ l of distilled water. Final Anti-MARCH2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at -20°C. Avoid repeat freeze-thaw cycles.
Precautions	MARCH2 Antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	MARCHF2 (HGNC:28038)
Synonyms	MARCH2, RNF172
Function	E3 ubiquitin-protein ligase that may mediate ubiquitination of TFRC and CD86, and promote their subsequent endocytosis and sorting to lysosomes via multivesicular bodies. E3 ubiquitin ligases accept ubiquitin from an E2 ubiquitin-conjugating enzyme in the form of a thioester and then directly transfer the ubiquitin to targeted substrates (PubMed: 14722266 , PubMed: 16428329). Together with GOPC/CAL mediates the ubiquitination and lysosomal degradation of CFTR (PubMed: 23818989). Ubiquitinates and

therefore mediates the degradation of DLG1 (PubMed:[17980554](#)). Regulates the intracellular trafficking and secretion of alpha1-antitrypsin/SERPINA1 and HP/haptoglobin via ubiquitination and degradation of the cargo receptor ERGIC3 (PubMed:[31142615](#)). Negatively regulates the antiviral and antibacterial immune response by repression of the NF-kB and type 1 IFN signaling pathways, via MARCHF2-mediated K48-linked polyubiquitination of IKBKG/NEMO, resulting in its proteasomal degradation (PubMed:[32935379](#)). May be involved in endosomal trafficking through interaction with STX6 (PubMed:[15689499](#)).


Cellular Location

Endoplasmic reticulum membrane; Multi-pass membrane protein {ECO:0000250|UniProtKB:Q5I0I2}. Lysosome membrane; Multi-pass membrane protein. Endosome membrane; Multi-pass membrane protein {ECO:0000250|UniProtKB:Q5I0I2}. Golgi apparatus membrane; Multi-pass membrane protein. Cytoplasm. Cell membrane; Multi-pass membrane protein

Tissue Location

Broadly expressed..

Images



Host: Rabbit
Target Name: MARCH2
Sample Tissue: Esophagus Tumor lysates
Antibody Dilution: 1.0µg/ml

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.