

RNF139 antibody - N-terminal region

Rabbit Polyclonal Antibody

Catalog # AI12216

Product Information

Application	WB
Primary Accession	Q8WU17
Other Accession	NM_007218 , NP_009149
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Dog, Guinea Pig, Horse, Bovine
Predicted	Rat, Rabbit, Pig, Dog, Guinea Pig, Horse, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	75994

Additional Information

Gene ID	11236
Alias Symbol	HRCA1, MGC31961, RCA1, TRC8
Other Names	E3 ubiquitin-protein ligase RNF139, 6.3.2.-, RING finger protein 139, Translocation in renal carcinoma on chromosome 8 protein, RNF139 (HGNC:17023)
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 50 ul of distilled water. Final anti-RNF139 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
Precautions	RNF139 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	RNF139 (HGNC:17023)
Function	E3-ubiquitin ligase; acts as a negative regulator of cell proliferation through mechanisms involving G2/M arrest and cell death (PubMed: 10500182 , PubMed: 12032852 , PubMed: 17016439). Required for MHC class I ubiquitination in cells expressing the cytomegalovirus protein US2 before dislocation from the endoplasmic reticulum (ER) (PubMed: 19720873). Affects SREBP processing by hindering the SREBP-SCAP complex translocation from the ER to the Golgi, thereby reducing SREBF2 target gene expression (PubMed: 19706601 , PubMed: 20068067). Involved in the sterol-accelerated degradation of HMGCR (PubMed: 22143767 , PubMed: 23223569). This is

achieved through binding of RNF139 to INSIG1 and/or INSIG2 at the ER membrane (PubMed:[22143767](#)). In addition, interaction of RNF139 with AUP1 facilitates interaction of RNF139 with ubiquitin-conjugating enzyme UBE2G2 and ubiquitin ligase AMFR, leading to ubiquitination of HMGCR (PubMed:[23223569](#)). The ubiquitinated HMGCR is then released from the ER into the cytosol for subsequent destruction (PubMed:[22143767](#), PubMed:[23223569](#)). Required for INSIG1 ubiquitination (PubMed:[20068067](#)). May be required for EIF3 complex ubiquitination (PubMed:[20068067](#)).

Cellular Location

Endoplasmic reticulum membrane; Multi-pass membrane protein

Tissue Location

Highly expressed in testis, placenta and adrenal gland. Moderate expression in heart, brain, liver, skeletal muscle and pancreas, and low expression in lung and kidney

Images



WB Suggested Anti-RNF139 Antibody Titration: 0.2-1
µg/ml

ELISA Titer: 1:312500

Positive Control: MCF7 cell lysate

RNF139 is supported by BioGPS gene expression data to
be expressed in MCF7

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.