

# RNF121 antibody - middle region

Rabbit Polyclonal Antibody Catalog # AI12285

#### **Product Information**

Application WB
Primary Accession Q9H920

Other Accession <u>NM 018320</u>, <u>NP 060790</u>

**Reactivity** Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Dog, Guinea Pig, Horse, Bovine

**Predicted** Human, Mouse, Rat, Pig, Chicken, Dog, Guinea Pig, Horse, Bovine

Host Rabbit
Clonality Polyclonal
Calculated MW 37882

## **Additional Information**

**Gene ID** 55298

Alias Symbol FLJ11099

Other Names RING finger protein 121, RNF121

Format Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium

azide and 2% sucrose.

**Reconstitution & Storage** Add 50 ul of distilled water. Final anti-RNF121 antibody concentration is 1

mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at

20°C. Avoid repeat freeze-thaw cycles.

**Precautions** RNF121 antibody - middle region is for research use only and not for use in

diagnostic or therapeutic procedures.

#### **Protein Information**

Name RNF121

**Function** E3 ubiquitin ligase which accepts ubiquitin and transfers it to substrates

thereby promoting their degradation by the endoplasmic

reticulum-associated degradation (ERAD) pathway which is a pathway involved in ubiquitin-dependent degradation of misfolded endoplasmic reticulum proteins (By similarity). May regulate the unfolded protein response

to reduce endoplasmic reticulum stress (By similarity).


**Cellular Location** Endoplasmic reticulum membrane {ECO:0000250|UniProtKB:Q09251};

Multi-pass membrane protein

### References

Gerhard, D.S., Genome Res. 14(10B), 2121-2127 (2004) Reconstitution and Storage: For short termuse, store at 2-8Cu pto 1 week. For long terms to rage, store at 2-20Cinsmall aliquots to prevent freeze-thaw cycles.

# **Images**



WB Suggested Anti-RNF121 Antibody Titration: 0.2-1

μg/ml

ELISA Titer: 1:2500

Positive Control: Human Thymus

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.