

RNF169 antibody - N-terminal region

Rabbit Polyclonal Antibody

Catalog # AI12287

Product Information

Application	WB
Primary Accession	Q8NCN4
Other Accession	NM_001098638 , NP_001092108
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Dog, Guinea Pig, Horse, Bovine
Predicted	Rabbit, Pig, Chicken, Dog, Guinea Pig, Horse, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	77194

Additional Information

Gene ID	254225
Alias Symbol	KIAA1991
Other Names	E3 ubiquitin-protein ligase RNF169, 6.3.2.-, RING finger protein 169, RNF169, KIAA1991
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 50 ul of distilled water. Final anti-RNF169 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
Precautions	RNF169 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	RNF169
Synonyms	KIAA1991
Function	Probable E3 ubiquitin-protein ligase that acts as a regulator of double-strand breaks (DSBs) repair following DNA damage. Functions in a non-canonical fashion to harness RNF168-mediated protein recruitment to DSB-containing chromatin, thereby contributing to regulation of DSB repair pathway utilization (PubMed: 22492721 , PubMed: 30773093). Once recruited to DSB repair sites by recognizing and binding ubiquitin catalyzed by RNF168, competes with TP53BP1 and BRCA1 for association with RNF168-modified chromatin, thereby favouring homologous recombination repair (HRR) and

single-strand annealing (SSA) instead of non-homologous end joining (NHEJ) mediated by TP53BP1 (PubMed:[30104380](#), PubMed:[30773093](#)). E3 ubiquitin-protein ligase activity is not required for regulation of DSBs repair.


Cellular Location

Chromosome. Nucleus, nucleoplasm. Note=Localizes to sites of double-strand breaks (DSBs) following DNA damage. Recruited to DSBs via recognition of RNF168-dependent ubiquitin products.

References

Olsen, J.V., (2006) Cell 127(3), 635-648 Reconstitution and Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.

Images



WB Suggested Anti-RNF169 Antibody Titration: 0.2-1 µg/ml
ELISA Titer: 1:312500
Positive Control: THP-1 cell lysate

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.