

Slc25a27 antibody - C-terminal region

Rabbit Polyclonal Antibody Catalog # AI12339

Product Information

Application WB
Primary Accession Q9D6D0

Other Accession <u>NM 028711, NP 082987</u>

Reactivity Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Dog, Guinea Pig, Horse, Bovine,

Sheep

Predicted Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Dog, Bovine

HostRabbitClonalityPolyclonalCalculated MW35798

Additional Information

Gene ID 74011

Alias Symbol 3632410G24Rik, 9430092A03Rik, D530043E16Rik, Ucp4

Format Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium

azide and 2% sucrose.

Reconstitution & Storage Add 50 ul of distilled water. Final anti-Slc25a27 antibody concentration is 1

mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at

20°C. Avoid repeat freeze-thaw cycles.

Precautions Slc25a27 antibody - C-terminal region is for research use only and not for use

in diagnostic or therapeutic procedures.

Protein Information

Name Slc25a27 {ECO:0000312 | MGI:MGI:1921261}

Function Facilitates proton transport across the inner mitochondrial membrane and


may dissipate excessive proton gradient associated with oxidative and metabolic stress at neuronal synapses. Regulates glutamate-induced proton conductance in astrocytes, shifting the energy metabolism toward aerobic glycolysis and lactate transfer to neurons for ATP synthesis. Can transport chloride ions with lower efficiency. The transport mechanism remains to be

elucidated.

Cellular Location Mitochondrion inner membrane; Multi-pass membrane protein. Cell

projection, neuron projection. Note=Localizes to neuronal cell body and processes. Within mitochondrial inner membrane, it is mainly observed in the inner boundary membrane locally separated from F(1)F(0) ATP synthase,

Images



WB Suggested Anti-Slc25a27 Antibody Titration: 1.0 $\mu g/ml$ Positive Control: Mouse Brain

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.