

Ppic Antibody - C-terminal region

Rabbit Polyclonal Antibody

Catalog # AI12412

Product Information

Application	WB
Primary Accession	P30412
Other Accession	NM_008908 , NP_032934
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Dog, Guinea Pig, Horse, Bovine
Predicted	Human, Mouse, Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	22794


Additional Information

Gene ID	19038
Alias Symbol	CyP-20°C
Other Names	Peptidyl-prolyl cis-trans isomerase C, PPIase C, 5.2.1.8, Cyclophilin C, Rotamase C, Ppic, Cypc
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 50 ul of distilled water. Final anti-Ppic antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
Precautions	Ppic Antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	Ppic {ECO:0000312 MGI:MGI:97751}
Function	PPIase that catalyzes the cis-trans isomerization of proline imidic peptide bonds in oligopeptides and may therefore assist protein folding.
Cellular Location	Cytoplasm.
Tissue Location	Specifically expressed in kidney.

Images



WB Suggested Anti-Ppic Antibody Titration: 1.0 µg/ml
Positive Control: Mouse Heart

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.