

# AWAT1 antibody - N-terminal region

Rabbit Polyclonal Antibody Catalog # AI12469

#### **Product Information**

Application WB
Primary Accession Q58HT5

Other Accession NM 001013579, NP 001013597

**Reactivity** Human, Mouse, Rat, Rabbit, Zebrafish, Dog, Guinea Pig, Horse, Bovine

**Predicted** Human, Mouse, Rat, Dog, Guinea Pig, Bovine

Host Rabbit
Clonality Polyclonal
Calculated MW 37759

### **Additional Information**

**Gene ID** 158833

Alias Symbol AWAT1, DGA2, DGAT2L3

**Other Names** Acyl-CoA wax alcohol acyltransferase 1, 2.3.1.75, Diacylglycerol

O-acyltransferase 2-like protein 3, Diacylglycerol acyltransferase 2, Long-chain-alcohol O-fatty-acyltransferase 1, AWAT1, DGA2, DGAT2L3

Format Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium

azide and 2% sucrose.

**Reconstitution & Storage** Add 50 ul of distilled water. Final anti-AWAT1 antibody concentration is 1

mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at

20°C. Avoid repeat freeze-thaw cycles.

**Precautions** AWAT1 antibody - N-terminal region is for research use only and not for use

in diagnostic or therapeutic procedures.

### **Protein Information**

Name AWAT1

**Synonyms** DGA2, DGAT2L3

**Function** Acyltransferase that catalyzes the formation of ester bonds between fatty

alcohols and fatty acyl-CoAs to form wax monoesters (PubMed: 15671038). Shows a strong preference for decyl alcohol (C10), with less activity towards C16 and C18 alcohols (PubMed: 15671038). Shows a strong preference for

saturated acyl-CoAs (PubMed: 15671038).


**Cellular Location** Endoplasmic reticulum membrane {ECO:0000250 | UniProtKB:Q6E213};

Multi-pass membrane protein

### **Tissue Location**

Predominantly expressed in skin, where it is limited to the sebaceous gland. Expressed in more mature, centrally located cells just before their rupture and sebum release. Also expressed in all tissues except spleen. Expressed at higher level in thymus, prostate and testis.

## **Images**



WB Suggested Anti-AWAT1 Antibody Titration: 0.2-1  $\mu g/ml$  Positive Control: Human Spleen

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.