

MPG antibody - C-terminal region

Rabbit Polyclonal Antibody

Catalog # AI12480

Product Information

Application	WB
Primary Accession	Q5J9I4
Other Accession	NM_001015052 , NP_001015052
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Dog, Guinea Pig, Horse, Bovine
Predicted	Human, Mouse, Rat, Rabbit, Pig, Guinea Pig, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	32 KDa

Additional Information

Alias Symbol	AAG, APNG, CRA36.1, MDG, Mid1, PIG11, PIG16, anpg, ADPG
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 50 ul of distilled water. Final anti-MPG antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
Precautions	MPG antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.



Protein Information

References

Oh,J.H.,(2005)Mamm.Genome16(12),942-954ReconstitutionandStorage:Forshorttermuse,storeat2-8Cupto1week.Forlongtermstorage,storeat-20Cinsmallaliquotstopreventfreeze-thawcycles.

Images

WB Suggested Anti-MPG Antibody Titration: 0.2-1 µg/ml
Positive Control: Jurkat cell lysate



WB Suggested Anti-MPG Antibody

Positive Control: Lane1: 50ug RCC4 lysate, Lane2: 50ug RCC4 lysate, Lane3: 50ug 786-0 lysate, Lane4: 50ug 786-0 lysate

Primary Antibody Dilution : 1:1000

Secondary Antibody : Anti-rabbit-Alexa 680

Secondary Antibody Dilution : 1:5000

Submitted by: Dr Ester Hammond, Gray Institute for Radiation Oncology and Biology-Department of Oncology, University of Oxford

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.