

FURIN antibody - N-terminal region

Rabbit Polyclonal Antibody

Catalog # AI12564

Product Information

Application	WB
Primary Accession	P09958
Other Accession	NM_002569 , NP_002560
Reactivity	Human, Mouse, Rat, Rabbit, Zebrafish, Dog, Guinea Pig, Horse, Bovine, Sheep
Predicted	Human, Mouse, Rat, Rabbit, Zebrafish, Chicken, Dog, Horse, Bovine, Sheep
Host	Rabbit
Clonality	Polyclonal
Calculated MW	86678

Additional Information

Gene ID	5045
Alias Symbol	FUR, PACE, PCSK3, SPC1
Other Names	Furin, 3.4.21.75, Dibasic-processing enzyme, Paired basic amino acid residue-cleaving enzyme, PACE, FURIN, FUR, PACE, PCSK3
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 50 ul of distilled water. Final anti-FURIN antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
Precautions	FURIN antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	FURIN {ECO:0000303 PubMed:7690548, ECO:0000312 HGNC:HGNC:8568}
Function	Ubiquitous endoprotease within constitutive secretory pathways capable of cleavage at the RX(K/R)R consensus motif (PubMed: 11799113 , PubMed: 1629222 , PubMed: 1713771 , PubMed: 2251280 , PubMed: 24666235 , PubMed: 25974265 , PubMed: 7592877 , PubMed: 7690548 , PubMed: 9130696). Mediates processing of TGFβ1, an essential step in TGF-beta-1 activation (PubMed: 7737999). Converts through proteolytic cleavage the non-functional Brain natriuretic factor prohormone into its active hormone BNP(1-32) (PubMed: 20489134 , PubMed: 21763278). By mediating processing of accessory subunit ATP6AP1/Ac45 of the V-ATPase, regulates the acidification of dense-core secretory granules in islets of Langerhans cells (By similarity).


Cellular Location

Golgi apparatus, trans-Golgi network membrane; Single-pass type I membrane protein. Cell membrane; Single-pass type I membrane protein. Secreted. Endosome membrane; Single-pass type I membrane protein. Note=Shuttles between the trans-Golgi network and the cell surface (PubMed:11799113, PubMed:9412467). Propeptide cleavage is a prerequisite for exit of furin molecules out of the endoplasmic reticulum (ER). A second cleavage within the propeptide occurs in the trans Golgi network (TGN), followed by the release of the propeptide and the activation of furin (PubMed:11799113)

Tissue Location

Seems to be expressed ubiquitously.

Images



WB Suggested Anti-FURIN Antibody Titration: 0.2-1 µg/ml
ELISA Titer: 1:62500
Positive Control: THP-1 cell lysate

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.