10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Fscn1 antibody - N-terminal region

Rabbit Polyclonal Antibody
Catalog # AI12653

Product Information

Application	WB
Primary Accession	Q61553
Other Accession	NM_007984, NP_032010
Reactivity	Human, Mouse, Rat, Zebrafish, Pig, Dog, Guinea Pig, Horse, Bovine
Predicted	Human, Mouse, Rat, Zebrafish, Pig, Chicken, Dog, Guinea Pig, Horse, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	54508

Additional Information


Gene ID	14086
Alias Symbol	AI663989, Fan1, fascin-1
Other Names	Fascin, Singed-like protein, Fscn1, Fan1, Snl
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 50 ul of distilled water. Final anti-Fscn1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
Precautions	Fscn1 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	Fscn1
Synonyms	Fan1, Snl
Function	Actin-binding protein that contains 2 major actin binding sites (By similarity). Organizes filamentous actin into parallel bundles (PubMed:7738015). Plays a role in the organization of actin filament bundles and the formation of microspikes, membrane ruffles, and stress fibers (By similarity). Important for the formation of a diverse set of cell protrusions, such as filopodia, and for cell motility and migration (PubMed:21685497). Mediates reorganization of the actin cytoskeleton and axon growth cone collapse in response to NGF (By similarity).

Cellular Location	Cytoplasm, cytosol {ECO:0000250|UniProtKB:Q16658}. Cytoplasm, cell cortex {ECO:0000250|UniProtKB:Q16658}. Cytoplasm, cytoskeleton {ECO:0000250|UniProtKB:Q16658}. Cytoplasm, cytoskeleton, stress fiber {ECO:0000250|UniProtKB:Q16658}. Cell projection, growth cone. Cell projection, filopodium. Cell projection, invadopodium. Cell projection, microvillus {ECO:0000250|UniProtKB:Q16658}. Cell junction {ECO:0000250|UniProtKB:Q16658}. Note=Colocalized with RUFY3 and F-actin at filipodia of the axonal growth cone (PubMed:24720729). Colocalized with DBN1 and F-actin at the transitional domain of the axonal growth cone (PubMed:24720729). {ECO:0000250|UniProtKB:Q16658, ECO:0000269|PubMed:24720729}
Tissue Location	Most abundant in brain. Detected at lower levels in lung, uterus, small intestine and spleen (at protein level)

Images



WB Suggested Anti-Fscn1 Antibody Titration: 1.0 µg/ml
Positive Control: Mouse Thymus

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.