

# SHMT2 antibody - N-terminal region

Rabbit Polyclonal Antibody

Catalog # AI12655

## Product Information

---

<b>Application</b>	WB, IHC
<b>Primary Accession</b>	<a href="#">P34897</a>
<b>Other Accession</b>	<a href="#">NM_005412</a> , <a href="#">NP_005403</a>
<b>Reactivity</b>	Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Dog, Guinea Pig, Horse, Bovine
<b>Predicted</b>	Human, Mouse, Rat, Zebrafish, Horse
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Calculated MW</b>	55993

## Additional Information

---

<b>Gene ID</b>	6472
<b>Alias Symbol</b>	GLYA, SHMT
<b>Other Names</b>	Serine hydroxymethyltransferase, mitochondrial, SHMT, 2.1.2.1, Glycine hydroxymethyltransferase, Serine methylase, SHMT2
<b>Format</b>	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
<b>Reconstitution &amp; Storage</b>	Add 100 ul of distilled water. Final anti-SHMT2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
<b>Precautions</b>	SHMT2 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	SHMT2 ( <a href="#">HGNC:10852</a> )
<b>Function</b>	Catalyzes the cleavage of serine to glycine accompanied with the production of 5,10-methylenetetrahydrofolate, an essential intermediate for purine biosynthesis (PubMed: <a href="#">24075985</a> , PubMed: <a href="#">25619277</a> , PubMed: <a href="#">29364879</a> , PubMed: <a href="#">33015733</a> ). Serine provides the major source of folate one-carbon in cells by catalyzing the transfer of one carbon from serine to tetrahydrofolate (PubMed: <a href="#">25619277</a> ). Contributes to the de novo mitochondrial thymidylate biosynthesis pathway via its role in glycine and tetrahydrofolate metabolism: thymidylate biosynthesis is required to prevent uracil accumulation in mtDNA (PubMed: <a href="#">21876188</a> ). Also required for mitochondrial translation by producing 5,10- methylenetetrahydrofolate; 5,10-methylenetetrahydrofolate providing

methyl donors to produce the taurinomethyluridine base at the wobble position of some mitochondrial tRNAs (PubMed:[29364879](#), PubMed:[29452640](#)). Associates with mitochondrial DNA (PubMed:[18063578](#)). In addition to its role in mitochondria, also plays a role in the deubiquitination of target proteins as component of the BRISC complex: required for IFNAR1 deubiquitination by the BRISC complex (PubMed:[24075985](#)).

### Cellular Location

Mitochondrion. Mitochondrion matrix, mitochondrion nucleoid. Mitochondrion inner membrane. Cytoplasm Nucleus. Note=Mainly localizes in the mitochondrion. Also found in the cytoplasm and nucleus as part of the BRISC complex (PubMed:24075985).


### References

---


Fu,T.F.,(2001)Arch.Biochem.Biophys.393(1),42-50ReconstitutionandStorage:Forshorttermuse,storeat2-8Cupto1week.Forlongtermstorage,storeat-20Cinsmallaliquotstopreventfreeze-thawcycles.

### Images

---



Human Heart



WB Suggested Antibody Titration: 2.5 µg/ml  
Positive Control: HepG2

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.