

B4galt5 antibody - C-terminal region

Rabbit Polyclonal Antibody

Catalog # AI12691

Product Information

Application	WB
Primary Accession	Q9JMK0
Other Accession	NM_019835 , NP_062809
Reactivity	Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Dog, Guinea Pig, Horse, Bovine
Predicted	Mouse, Rat, Rabbit, Pig, Chicken, Dog, Guinea Pig, Horse, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	44814

Additional Information

Gene ID	56336
Alias Symbol	9430078I07Rik, AW049941, AW539721
Other Names	Beta-1, 4-galactosyltransferase 5, Beta-1, 4-GalTase 5, Beta4Gal-T5, b4Gal-T5, 2.4.1.-, Beta-1, 4-GalT II, UDP-Gal:beta-GlcNAc beta-1, 4-galactosyltransferase 5, UDP-galactose:beta-N-acetylglucosamine beta-1, 4-galactosyltransferase 5, B4galt5, Bgt-5
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 50 ul of distilled water. Final anti-B4galt5 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
Precautions	B4galt5 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	B4galt5 {ECO:0000312 MGI:MGI:1927169}
Synonyms	Bgt-5
Function	Catalyzes the synthesis of lactosylceramide (LacCer) via the transfer of galactose from UDP-galactose to glucosylceramide (GlcCer) (PubMed: 21057870 , PubMed: 23882130 , PubMed: 30114188). LacCer is the starting point in the biosynthesis of all gangliosides (membrane-bound glycosphingolipids) which play pivotal roles in the CNS including neuronal maturation and axonal and myelin formation (PubMed: 30114188). Plays a

role in the glycosylation of BMPR1A and regulation of its protein stability (PubMed:[29415997](#)). Essential for extraembryonic development during early embryogenesis (PubMed:[20574042](#), PubMed:[21057870](#)).


Cellular Location

Golgi apparatus, Golgi stack membrane {ECO:0000250|UniProtKB:P15291}; Single-pass type II membrane protein Golgi apparatus {ECO:0000250|UniProtKB:A0A1S6M251}. Note=Trans cisternae of Golgi stack. {ECO:0000250|UniProtKB:P15291}

Tissue Location

Highest levels in heart, brain, liver and kidney with lower levels in spleen, lung and testis

Images



WB Suggested Anti-B4galt5 Antibody Titration: 0.2-1
µg/ml
ELISA Titer: 1:62500
Positive Control: Mouse Kidney

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.