

IMPAD1 antibody - middle region

Rabbit Polyclonal Antibody

Catalog # AI12778

Product Information

Application	WB
Primary Accession	Q9NX62
Other Accession	NM_017813 , NP_060283
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Dog, Guinea Pig, Horse, Bovine
Predicted	Rat, Rabbit, Chicken, Dog, Guinea Pig, Horse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	38681

Additional Information

Gene ID	54928
Alias Symbol	FLJ20421, IMPA3, GPAPP, IMP 3, IMP-3
Other Names	Inositol monophosphatase 3, IMP 3, IMPase 3, 3.1.3.25, 3.1.3.7, Golgi 3-prime phosphoadenosine 5-prime phosphate 3-prime phosphatase, Golgi-resident PAP phosphatase, gPAPP, Inositol monophosphatase domain-containing protein 1, Inositol-1(or 4)-monophosphatase 3, Myo-inositol monophosphatase A3, IMPAD1, IMPA3
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 50 ul of distilled water. Final anti-IMPAD1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
Precautions	IMPAD1 antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	BPNT2 (HGNC:26019)
Synonyms	IMPA3, IMPAD1
Function	Exhibits 3'-nucleotidase activity toward adenosine 3',5'- bisphosphate (PAP), namely hydrolyzes adenosine 3',5'-bisphosphate into adenosine 5'-monophosphate (AMP) and a phosphate. May play a role in the formation of skeletal elements derived through endochondral ossification, possibly by clearing adenosine 3',5'-bisphosphate produced by Golgi sulfotransferases

during glycosaminoglycan sulfation. Has no activity toward 3'-phosphoadenosine 5'-phosphosulfate (PAPS) or inositol phosphate (IP) substrates including I(1)P, I(1,4)P2, I(1,3,4)P3, I(1,4,5)P3 and I(1,3,4,5)P4.


Cellular Location

Golgi apparatus. Golgi apparatus, trans-Golgi network membrane; Single-pass type II membrane protein {ECO:0000255, ECO:0000269 | PubMed:18695242}.
Note=The catalytic core is predicted to reside within the Golgi lumen

References

Zhang,H.,(2003)Nat.Biotechnol.21(6),660-666ReconstitutionandStorage:Forshorttermuse,storeat2-8Cupto1week.Forlongtermstorage,storeat-20Cinsmallaliquotstopreventfreeze-thawcycles.

Images



WB Suggested Anti-IMPAD1 Antibody Titration: 0.2-1
µg/ml
ELISA Titer: 1:312500
Positive Control: Human Spleen

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.