

Tnik antibody - middle region

Rabbit Polyclonal Antibody

Catalog # AI12939

Product Information

Application	WB
Primary Accession	P83510
Other Accession	NM_001163008 , NP_001156480
Reactivity	Human, Mouse, Rat, Rabbit, Zebrafish, Dog, Guinea Pig, Horse, Bovine, Yeast
Predicted	Human, Mouse, Rat, Rabbit, Chicken, Dog, Guinea Pig, Horse, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	150367

Additional Information

Gene ID	665113
Alias Symbol	1500031A17Rik, 4831440I19Rik, AI451411, C530008O15Rik, C630040K21Rik, KIAA0551, MGC189819, MGC189859
Other Names	Traf2 and NCK-interacting protein kinase, 2.7.11.1, Tnik, Kiaa0551
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 50 ul of distilled water. Final anti-Tnik antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
Precautions	Tnik antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information


Name	Tnik
Synonyms	Kiaa0551
Function	Serine/threonine kinase that acts as an essential activator of the Wnt signaling pathway. Recruited to promoters of Wnt target genes and required to activate their expression. May act by phosphorylating TCF4/TCF7L2. Appears to act upstream of the JUN N- terminal pathway. May play a role in the response to environmental stress. Part of a signaling complex composed of NEDD4, RAP2A and TNIK which regulates neuronal dendrite extension and arborization during development. More generally, it may play a role in cytoskeletal rearrangements and regulate cell spreading. Phosphorylates

SMAD1 on Thr-322. Activator of the Hippo signaling pathway which plays a pivotal role in organ size control and tumor suppression by restricting proliferation and promoting apoptosis. MAP4Ks act in parallel to and are partially redundant with STK3/MST2 and STK4/MST2 in the phosphorylation and activation of LATS1/2, and establish MAP4Ks as components of the expanded Hippo pathway.

Cellular Location

Nucleus. Cytoplasm. Recycling endosome Cytoplasm, cytoskeleton.
Note=Associated with recycling endosomes and the cytoskeletal fraction upon RAP2A overexpression

Images



WB Suggested Anti-Tnik Antibody Titration: 1.0 µg/ml
Positive Control: Mouse Spleen

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.