

PWWP2B antibody - N-terminal region

Rabbit Polyclonal Antibody Catalog # AI12971

Product Information

Application WB
Primary Accession Q6NUJ5

Other Accession NM 001098637, NP 001092107
Reactivity Human, Mouse, Rat, Pig, Bovine

Predicted Mouse, Rat, Bovine

Host Rabbit
Clonality Polyclonal
Calculated MW 63967

Additional Information

Gene ID 170394

Alias Symbol FLJ39621, FLJ46823, PWWP2, RP11-273H7.1, bA432J24.1, pp8607

Other Names PWWP domain-containing protein 2B, PWWP2B, PWWP2

Format Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium

azide and 2% sucrose.

Reconstitution & Storage Add 50 ul of distilled water. Final anti-PWWP2B antibody concentration is 1

mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at

20°C. Avoid repeat freeze-thaw cycles.

Precautions PWWP2B antibody - N-terminal region is for research use only and not for use

in diagnostic or therapeutic procedures.

Protein Information

Name PWWP2B

Synonyms PWWP2

Function Chromatin-binding protein that acts as an adapter between distinct


nucleosome components (H3K36me3 or H2A.Z) and chromatin- modifying complexes, contributing to the regulation of the levels of histone acetylation at actively transcribed genes (PubMed:30228260). Competes with CHD4 and MBD3 for interaction with MTA1 to form a NuRD subcomplex, preventing the formation of full NuRD complex (containing CHD4 and MBD3), leading to recruitment of HDACs to gene promoters resulting in turn in the deacetylation of nearby H3K27 and H2A.Z (PubMed:30228260). Plays a role in facilitating transcriptional elongation through regulation of histone acetylation (By

similarity). Negatively regulates brown adipocyte thermogenesis by interacting with and stabilizing HDAC1 at the UCP1 gene promoter, thereby promoting histone deacetylation at the promoter leading to the repression of UCP1 expression (By similarity).

References

Deloukas P., et al. Nature 429:375-381(2004). Ota T., et al. Nat. Genet. 36:40-45(2004).

Images



WB Suggested Anti-PWWP2B Antibody Titration: 0.2-1

μg/ml

ELISA Titer: 1:62500

Positive Control: DU145 cell lysate

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.