

Tceal7 antibody - middle region

Rabbit Polyclonal Antibody

Catalog # AI13229

Product Information

Application	WB
Primary Accession	A3KGA4
Other Accession	NM_001127169 , NP_001120641
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Dog, Horse, Bovine, Yeast
Predicted	Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	11697

Additional Information

Gene ID	100040972
Alias Symbol	1110018P05Rik, AI118095
Other Names	Transcription elongation factor A protein-like 7, TCEA-like protein 7, Transcription elongation factor S-II protein-like 7, Tceal7
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 50 ul of distilled water. Final anti-Tceal7 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
Precautions	Tceal7 antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.


Protein Information

Name	Tceal7
Function	Plays a role in the negative regulation of NF-kappa-B signaling at the basal level by modulating transcriptional activity of NF-kappa-B on its target gene promoters. Associates with cyclin D1 promoter containing Myc E-box sequence and transcriptionally represses cyclin D1 expression. Regulates telomerase reverse transcriptase expression and telomerase activity in both ALT (alternative lengthening of telomeres) and telomerase-positive cell lines (By similarity).
Cellular Location	Nucleus.

References

Church D.M.,et al.PLoS Biol. 7:E1000112-E1000112(2009).

Images



WB Suggested Anti-Tceal7 Antibody Titration: 1.0 µg/ml
Positive Control: Mouse Small Intestine

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.