

Vgll2 antibody - N-terminal region

Rabbit Polyclonal Antibody

Catalog # AI13232

Product Information

Application	WB
Primary Accession	Q8BGW8
Other Accession	NM_153786 , NP_722481
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Dog, Guinea Pig, Horse, Bovine
Predicted	Human, Mouse, Rat, Rabbit, Pig, Chicken, Dog, Guinea Pig, Horse, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	34064

Additional Information

Gene ID	215031
Alias Symbol	C130057C21Rik, VITO-1, Vito1
Other Names	Transcription cofactor vestigial-like protein 2, Vgl-2, Protein VITO1, Vgll2, Vgl2, Vito1
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 50 ul of distilled water. Final anti-Vgll2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
Precautions	Vgll2 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.


Protein Information

Name	Vgll2
Synonyms	Vgl2, Vito1
Function	May act as a specific coactivator for the mammalian TEFs. May play a role in the development of skeletal muscles.
Cellular Location	Nucleus.
Tissue Location	Skeletal muscle specific.

References

Maeda T.,et al.J. Biol. Chem. 277:48889-48898(2002).
Mielcarek M.,et al.Gene Expr. Patterns 2:305-310(2002).
Carninci P.,et al.Science 309:1559-1563(2005).

Images



WB Suggested Anti-Vgll2 Antibody Titration: 1.0 µg/ml
Positive Control: Mouse Small Intestine

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.