

UBTFL1 antibody - N-terminal region

Rabbit Polyclonal Antibody

Catalog # AI13264

Product Information

Application	WB
Primary Accession	P0CB47
Other Accession	XM_926094 , XP_931187
Reactivity	Human, Rat, Guinea Pig
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	46134

Additional Information

Gene ID	642623
Alias Symbol	HMGPI, C11orf27
Other Names	Putative upstream-binding factor 1-like protein 1, UBTFL1, C11orf27
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 50 ul of distilled water. Final anti-UBTFL1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
Precautions	UBTFL1 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	UBTFL1 (HGNC:14533)
Synonyms	C11orf27
Function	Essential for proliferation of the inner cell mass and trophectodermal cells in peri-implantation development.
Cellular Location	Cytoplasm {ECO:0000250 UniProtKB:Q3USZ2}. Nucleus {ECO:0000250 UniProtKB:Q3USZ2, ECO:0000255 PROSITE-ProRule:PRU00267} Note=Mainly cytoplasmic from the 4-cell stage to the morula stage Becomes nuclear at the blastocyst stage {ECO:0000250 UniProtKB:Q3USZ2}

Images



WB Suggested Anti-UBTFL1 Antibody Titration: 0.2-1
µg/ml
Positive Control: MCF7 cell lysate

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.